

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

---

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 39/395, C07K 16/00, 16/18, 16/28, 16/30</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/25543</b> <b>(43) International Publication Date:</b> 28 September 1995 (28.09.95)
<b>(21) International Application Number:</b> PCT/US95/03035 <b>(22) International Filing Date:</b> 9 March 1995 (09.03.95) <b>(30) Priority Data:</b> 08/210,715 18 March 1994 (18.03.94) US 08/366,665 30 December 1994 (30.12.94) US <b>(71) Applicant:</b> THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10666 North Torrey Pines Road, La Jolla, CA 92037 (US). <b>(72) Inventors:</b> BROOKS, Peter; Apartment 6113, 3550 Lebon Drive, San Diego, CA 92122 (US). CHERESH, David, A.; 2108 Sea Village Circle, Cardiff, CA 92007 (US). <b>(74) Agents:</b> FITTING, Thomas et al.; The Scripps Research Institute, 10666 North Torrey Pines Road, TPC-8, La Jolla, CA 92037 (US).	<b>(81) Designated States:</b> AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).  <b>Published</b> <i>With international search report.</i>	
<b>(54) Title:</b> METHODS AND COMPOSITIONS USEFUL FOR INHIBITION OF ANGIOGENESIS <b>(57) Abstract</b>  The present invention describes methods for inhibiting angiogenesis in tissues using vitronectin $\alpha_v\beta_3$ antagonists, and particularly for inhibiting angiogenesis in inflamed tissues and in tumor tissues and metastases using therapeutic compositions containing $\alpha_v\beta_3$ antagonists.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

METHODS AND COMPOSITIONS USEFUL FOR  
INHIBITION OF ANGIOGENESIS

Technical Field

5           The present invention relates generally to the  
field of medicine, and relates specifically to  
methods and compositions for inhibiting  
angiogenesis of tissues using antagonists of the  
vitronectin receptor  $\alpha_v\beta_3$ .

10           Background

Integrins are a class of cellular receptors  
known to bind extracellular matrix proteins, and  
therefore mediate cell-cell and cell-extracellular  
15           matrix interactions, referred generally to as cell  
adhesion events. However, although many integrins  
and the ligands that bind an integrin are described  
in the literature, the biological function of many  
of the integrins remains elusive. The integrin  
20           receptors constitute a family of proteins with  
shared structural characteristics of noncovalent  
heterodimeric glycoprotein complexes formed of  $\alpha$   
and  $\beta$  subunits.

25           The vitronectin receptor, named for its  
original characteristic of preferential binding to  
vitronectin, is now known to refer to three  
different integrins, designated  $\alpha_v\beta_1$ ,  $\alpha_v\beta_3$ , and  $\alpha_v\beta_5$ .  
Horton, Int. J. Exp. Pathol., 71:741-759 (1990).  
 $\alpha_v\beta_1$  binds fibronectin and vitronectin.  $\alpha_v\beta_3$  binds a  
30           large variety of ligands, including fibrin,  
fibrinogen, laminin, thrombospondin, vitronectin,  
von Willebrand's factor, osteospondin and bone  
sialoprotein I.  $\alpha_v\beta_5$  binds vitronectin. The  
specific cell adhesion roles these three integrins  
35           play in the many cellular interactions in tissues  
is still under investigation, but it is clear that  
there are different integrins with different  
biological functions.

- 2 -

One important recognition site in the ligand for many integrins is the arginine-glycine-aspartic acid (RGD) tripeptide sequence. RGD is found in all of the ligands identified above for the vitronectin receptor integrins. This RGD recognition site can be mimicked by polypeptides ("peptides") that contain the RGD sequence, and such RGD peptides are known inhibitors of integrin function. It is important to note, however, that depending upon the sequence and structure of the RGD peptide, the specificity of the inhibition can be altered to target specific integrins.

For discussions of the RGD recognition site, see Pierschbacher et al., Nature, 309:30-33 (1984), and Pierschbacher et al., Proc. Natl. Acad. Sci. USA, 81:5985-5988 (1984). Various RGD polypeptides of varying integrin specificity have also been described by Grant et al., Cell, 58:933-943 (1989), Cheresh, et al., Cell, 58:945-953 (1989), Aumailley et al., FEBS Letts., 291:50-54 (1991), and Pfaff et al., J. Biol. Chem., 269:20233-20238 (1994), and in United States Patent Nos. 4,517,686, 4,578,079, 4,589,881, 4,614,517, 4,661,111, 4,792,525, 4,683,291, 4,879,237, 4,988,621, 5,041,380 and 5,061,693.

Angiogenesis is a process of tissue vascularization that involves the growth of new developing blood vessels into a tissue, and is also referred to as neo-vascularization. The process is mediated by the infiltration of endothelial cells and smooth muscle cells. The process is believed to proceed in any one of three ways: the vessels can sprout from pre-existing vessels, de-novo development of vessels can arise from precursor cells (vasculogenesis), or existing small vessels can enlarge in diameter. Blood et al., Bioch. Biophys. Acta, 1032:89-118 (1990). Vascular endothelial cells are known to contain at least

- 3 -

five RGD-dependent integrins, including the vitronectin receptor ( $\alpha_v\beta_3$  or  $\alpha_v\beta_5$ ), the collagen Types I and IV receptor ( $\alpha_1\beta_1$ ), the laminin receptor ( $\alpha_2\beta_1$ ), the fibronectin/laminin/collagen receptor ( $\alpha_3\beta_1$ ) and the fibronectin receptor ( $\alpha_5\beta_1$ ). Davis et al., J. Cell. Biochem., 51:206-218 (1993). The smooth muscle cell is known to contain at least six RGD-dependent integrins, including  $\alpha_5\beta_1$ ,  $\alpha_v\beta_3$  and  $\alpha_v\beta_5$ .

Angiogenesis is an important process in neonatal growth, but is also important in wound healing and in the pathogenesis of a large variety of clinical diseases including tissue inflammation, arthritis, tumor growth, diabetic retinopathy, macular degeneration by neovascularization of retina and the like conditions. These clinical manifestations associated with angiogenesis are referred to as angiogenic diseases. Folkman et al., Science, 235:442-447 (1987). Angiogenesis is generally absent in adult or mature tissues, although it does occur in wound healing and in the corpus leuteum growth cycle. See, for example, Moses et al., Science, 248:1408-1410 (1990).

It has been proposed that inhibition of angiogenesis would be a useful therapy for restricting tumor growth. Inhibition of angiogenesis has been proposed by (1) inhibition of release of "angiogenic molecules" such as  $\beta$ FGF (fibroblast growth factor), (2) neutralization of angiogenic molecules, such as by use of anti- $\beta$ FGF antibodies, and (3) inhibition of endothelial cell response to angiogenic stimuli. This latter strategy has received attention, and Folkman et al., Cancer Biology, 3:89-96 (1992), have described several endothelial cell response inhibitors, including collagenase inhibitor, basement membrane turnover inhibitors, angiostatic steroids, fungal-derived angiogenesis inhibitors, platelet factor 4,

- 4 -

thrombospondin, arthritis drugs such as D-penicillamine and gold thiomalate, vitamin D<sub>3</sub> analogs, alpha-interferon, and the like that might be used to inhibit angiogenesis. For additional proposed inhibitors of angiogenesis, see Blood et al., Bioch. Biophys. Acta., 1032:89-118 (1990), Moses et al., Science, 248:1408-1410 (1990), Ingber et al., Lab. Invest., 59:44-51 (1988), and United States Patent Nos. 5,092,885, 5,112,946, 5,192,744, and 5,202,352. None of the inhibitors of angiogenesis described in the foregoing references are targeted at inhibition of  $\alpha_v\beta_3$ .

RGD-containing peptides that inhibit vitronectin receptor  $\alpha_v\beta_3$  have also been described. Aumailley et al., FEBS Letts., 291:50-54 (1991), Choi et al., J. Vasc. Surg., 19:125-134 (1994), Smith et al., J. Biol. Chem., 265:12267-12271 (1990), and Pfaff et al., J. Biol. Chem., 269:20233-20238 (1994). However, the role of the integrin  $\alpha_v\beta_3$  in angiogenesis has never been suggested nor identified until the present invention.

Inhibition of cell adhesion in vitro using monoclonal antibodies immunospecific for various integrin  $\alpha$  or  $\beta$  subunits have implicated  $\alpha_v\beta_3$  in cell adhesion of a variety of cell types including microvascular endothelial cells. Davis et al., J. Cell. Biol., 51:206-218 (1993). In addition, Nicosia et al., Am. J. Pathol., 138:829-833 (1991), described the use of the RGD peptide GRGDS to in vitro inhibit the formation of "microvessels" from rat aorta cultured in collagen gel. However, the inhibition of formation of "microvessels" in vitro in collagen gel cultures is not a model for inhibition of angiogenesis in a tissue because it is not shown that the microvessel structures are the same as capillary sprouts or that the formation of the microvessel in collagen gel culture is the

- 5 -

same as neo-vascular growth into an intact tissue, such as arthritic tissue, tumor tissue or disease tissue where inhibition of angiogenesis is desirable.

Therefore, other than the studies reported here, Applicants are unaware of any other demonstration that angiogenesis could be inhibited in a tissue using inhibitors of cell adhesion. In particular, it has never been previously demonstrated that  $\alpha_v\beta_3$  function is required for angiogenesis in a tissue or that  $\alpha_v\beta_3$  antagonists can inhibit angiogenesis in a tissue.

#### Brief Description of the Invention

The present invention disclosure demonstrates that angiogenesis in tissues requires integrin  $\alpha_v\beta_3$ , and that inhibitors of  $\alpha_v\beta_3$  can inhibit angiogenesis. The disclosure also demonstrates that antagonists of other integrins, such as  $\alpha_v\beta_5$ , or  $\alpha_v\beta_1$ , do not inhibit angiogenesis, presumably because these other integrins are not essential for angiogenesis to occur.

The invention therefore describes methods for inhibiting angiogenesis in a tissue comprising administering to the tissue a composition comprising an angiogenesis-inhibiting amount of an  $\alpha_v\beta_3$  antagonist.

The tissue to be treated can be any tissue in which inhibition of angiogenesis is desirable, such as diseased tissue where neo-vascularization is occurring. Exemplary tissues include inflamed tissue, solid tumors, metastases, tissues undergoing restenosis, and the like tissues.

An  $\alpha_v\beta_3$  antagonist for use in the present methods is capable of binding to  $\alpha_v\beta_3$  and competitively inhibiting the ability of  $\alpha_v\beta_3$  to bind to a natural ligand. Preferably, the antagonist exhibits specificity for  $\alpha_v\beta_3$  over other integrins.



- 6 -

In a particularly preferred embodiment, the  $\alpha_v\beta_3$  antagonist inhibits binding of fibrinogen or other RGD-containing ligands to  $\alpha_v\beta_3$ , but does not substantially inhibit binding of fibronectin to  $\alpha_{IIB}\beta_3$ . A preferred  $\alpha_v\beta_3$  antagonist can be a polypeptide or a monoclonal antibody, or functional fragment thereof, that immunoreacts with  $\alpha_v\beta_3$ .

#### Brief Description of the Drawings

In the drawings forming a portion of this disclosure:

Figures 1A-1D illustrate the tissue distribution of the integrin subunits,  $\beta_3$  and  $\beta_1$ , in normal skin and in skin undergoing wound healing designated as granulation tissue.

Immunohistochemistry with antibodies to  $\beta_3$  and  $\beta_1$  was performed as described in Example 3A. Figures 1A and 1B respectively illustrate the immunoreactivity of anti- $\beta_3$  in normal skin and granulation tissue. Figures 1C and 1D respectively illustrate the immunoreactivity of anti- $\beta_1$  in normal skin and granulation tissue.

Figures 2A-2D illustrate the tissue distribution of the von Willebrand factor and laminin ligands that respectively bind the  $\beta_3$  and  $\beta_1$  integrin subunits in normal skin and in skin undergoing wound healing designated as granulation tissue. Immunohistochemistry with antibodies to von Willebrand factor (anti-vWF) and laminin (anti-laminin) was performed as described in Example 3B. Figures 2A and 2B respectively illustrate the immunoreactivity of anti-vWF in normal skin and granulation tissue. Figures 2C and 2D respectively illustrate the immunoreactivity of anti-laminin in normal skin and granulation tissue.

Figures 3A-3D illustrate the tissue distribution of the vitronectin integrin receptor,  $\alpha_v\beta_3$ , in tissue biopsies of bladder cancer, colon

- 7 -

cancer, breast cancer and lung cancer, respectively. Immunohistochemistry with the LM609 antibody against  $\alpha_v\beta_3$  was performed as described in Example 3C.

5           Figure 4 illustrates a typical photomicrograph of a CAM of this invention devoid of blood vessels in an untreated 10 day old chick embryo. The preparation is described in Example 5B.

10           Figures 5A-5C illustrate the tissue distribution of the integrins  $\beta_1$  and  $\alpha_v\beta_3$  in the CAM preparation of this invention. Figure 5A shows the distribution of the  $\beta_1$  subunit in an untreated 10 day old CAM preparation as detected by immunofluorescence immunoreactivity with CSAT, an anti- $\beta_1$  antibody. Figure 5B shows the distribution of the  $\alpha_v\beta_3$  receptor in an untreated 10 day old CAM preparation as detected by immunofluorescence immunoreactivity with LM609, an anti- $\alpha_v\beta_3$  antibody. Figure 5C shows the distribution of the  $\alpha_v\beta_3$  receptor in an  $\beta$ FGF treated 10 day old CAM preparation as detected by immunofluorescence immunoreactivity with LM609, an anti- $\alpha_v\beta_3$  antibody. The treatments and results are described in Example 5C.

25           Figure 6 illustrates the quantification in a bar graph of the relative expression of  $\alpha_v\beta_3$  and  $\beta_1$  in untreated and  $\beta$ FGF treated 10 day old CAMs as described in Example 6A. The mean fluorescence intensity is plotted on the Y-axis with the integrin profiles plotted on the X-axis.

30           Figures 7A-7C illustrates the appearance of an untreated 10 day old CAM, a  $\beta$ FGF treated CAM, and a  $\text{TNF}\alpha$  treated CAM, respectively, the procedures and results of which are described in Example 6A.

35           Figures 8A-8E illustrate the effect of topical antibody treatment on FGF-induced angiogenesis in a day 10 CAM as described in Example 7A1). Figure 8A shows an untreated CAM preparation that is devoid

- 8 -

of blood vessels. Figure 8B shows the infiltration of new vasculature into an area previously devoid of vasculature induced by  $\beta$ FGF treatment. Figures 8C, 8D and 8E respectively show the effects of antibodies against  $\beta_1$  (anti- $\beta_1$ ; CSAT),  $\alpha_v\beta_5$  (anti- $\alpha_v\beta_5$ ; P3G2) and  $\alpha_v\beta_3$  (anti- $\alpha_v\beta_3$ ; LM609).

Figures 9A-9C illustrate the effect of intravenous injection of synthetic peptide 66203 on angiogenesis induced by tumors as described in Example 7D2). Figure 9A shows the lack of inhibitory effect of intravenous treatment with a control peptide (control peptide tumor) on angiogenesis resulting from tumor induction. The inhibition of such angiogenesis by intravenous injection of peptide 66203 (cyclic RGD tumor) is shown in Figure 9B. The lack of inhibitory effects or cytotoxicity on mature preexisting vessels following intravenous infusion of peptide 66203 in an area adjacent to the tumor-treated area is shown in Figure 9C (cyclic RGD adjacent CAM).

Figures 10A-10C illustrate the effect of intravenous application of monoclonal antibodies to growth factor induced angiogenesis as described in Example 7B1). Figure 10A shows  $\beta$ FGF-induced angiogenesis not exposed to antibody treatment (control). No inhibition of angiogenesis resulted when a similar preparation was treated with anti- $\alpha_v\beta_5$  antibody P3G2 as shown in Figure 10B. Inhibition of angiogenesis resulted with treatment of anti- $\alpha_v\beta_3$  antibody LM609 as shown in Figure 10C.

Figures 11A-11C illustrate the effect on embryonic angiogenesis following topical application of anti-integrin antibodies as described in Example 7C. Angiogenesis was not inhibited by treatment of a 6 day CAM with anti- $\beta_1$  and anti- $\alpha_v\beta_5$  antibodies respectively shown in Figures 11A and 11B. In contrast, treatment with the anti- $\alpha_v\beta_3$  antibody LM609 resulted in the

- 9 -

inhibition of blood vessel formation as shown in Figure 11C.

Figure 12 illustrates the quantification of the number of vessels entering a tumor in a CAM preparation as described in Example 7D1). The graph shows the number of vessels as plotted on the Y-axis resulting from topical application of either CSAT (anti- $\beta_1$ ), LM609 (anti- $\alpha_v\beta_3$ ) or P3G2 (anti- $\alpha_v\beta_5$ ).

Figures 13A-13D illustrate a comparison between wet tumor weights 7 days following treatment and initial tumor weights as described in Example 9A1)a. Each bar represents the mean  $\pm$  S.E. of 5-10 tumors per group. Tumors were derived from human melanoma (M21-L) (Figure 13A), pancreatic carcinoma (Fg) (Figure 13B), lung carcinoma (UCLAP-3) (Figure 13C), and laryngeal carcinoma (HEp3) (Figure 13D) CAM preparations and treated intravenously with PBS, CSAT (anti- $\beta_1$ ), or LM609 (anti- $\alpha_v\beta_3$ ). The graphs show the tumor weight as plotted on the Y-axis resulting from intravenous application of either CSAT (anti- $\beta_1$ ), LM609 (anti- $\alpha_v\beta_3$ ) or PBS as indicated on the X-axis.

Figures 14A and 14B illustrate histological sections of tumors treated with the P3G2 (anti- $\alpha_v\beta_5$ ) (Figure 14A) and LM609 (anti- $\alpha_v\beta_3$ ) (Figure 14B), and stained with hematoxylin and eosin as described in Example 9A1)a. As shown in Figure 14A, tumors treated with control antibody (P3G2) showed numerous viable and actively dividing tumor cells as indicated by mitotic figures (arrowheads) as well as by multiple blood vessels (arrows) throughout the tumor stroma. In contrast, few if any viable tumor cells or blood vessels were detected in tumors treated with LM609 (anti- $\alpha_v\beta_3$ ) in Figure 14B.

Figures 15A-15E correspond to M21L tumors treated with peptides as described in Example 9A1)b

- 10 -

and are as follows: Figure 15A, control cyclic RAD peptide (69601); Figure 15B, cyclic RGD peptide (66203); Figure 15C, adjacent CAM tissue taken from the same embryos treated with cyclic RGD peptide (66203) and high magnification (13x) of tumors treated with the control RAD (69601) in Figure 15D or cyclic RGD peptide (66203) in Figure 15E. Figure 15D depicts normal vessels from the RAD control peptide (69601) treated tumor. Figure 15E depicts examples of disrupted blood vessels from cyclic RGD peptide (66203) treated tumors (arrows).

Figures 16A-16E represent inhibition of angiogenesis by antagonists of angiogenesis in the in vivo rabbit eye model assay as described in Example 10. Figure 16A and 16B depict angiogenesis of the rabbit eye in the presence of  $\beta$ FGF and mAb P1F6 (anti- $\alpha_v\beta_5$ ). Figure 16C, 16D and 16E depict inhibition of angiogenesis of the rabbit eye in the presence of  $\beta$ FGF and mAb LM609 (anti- $\alpha_v\beta_3$ ).

Figure 17 represents a flow chart of how the in vivo mouse:human chimeric mouse model was generated as described in Example 11A. A portion of skin from a SCID mouse was replaced with human neonatal foreskin and allowed to heal for 4 weeks. After the graft had healed, the human foreskin was inoculated with human tumor cells. During the following 4 week period, a measurable tumor was established which comprised a human tumor with human vasculature growing from the human skin into the human tumor.

Figure 18 illustrates the percent of single cells derived from mAb-treated and peptide-treated CAMs and stained with Apop Tag as determined by FACS analysis and described in Examples 12A and 12B, respectively. The black and stippled bars represent cells from embryos treated 24 hours and 48 hours prior to the assay, respectively. Each

- 11 -

bar represents the mean  $\pm$  S.E. of three replicates. CAMs were treated mAb LM609 (anti- $\alpha_v\beta_3$ ), or CSAT (anti- $\beta_1$ ), or PBS as described in Example 12A2). CAMs were also treated with cyclic peptide 69203 (cyclo-RGDfV, indicated as Peptide 203) or control cyclic peptide 69601 (cyclo-RADfV, indicated as Peptide 601) as described in Example 12B.

Figures 19A and 19B illustrate the combined results of single cell suspensions of CAMs from embryos treated with either CSAT (anti- $\beta_1$ ) (Figure 19A) or LM609 (anti- $\alpha_v\beta_3$ ) (Figure 19B), stained with Apop Tag and propidium iodide, and analyzed by FACS as described in Example 12C. The Y axis represents Apop Tag staining in numbers of cells (Apoptosis), the X axis represents propidium iodide staining (DNA content). The horizontal line represents the negative gate for Apop Tag staining. The left and right panels indicates CAM cells from CSAT (anti- $\beta_1$ ) (Figure 19A) and LM609 (anti- $\alpha_v\beta_3$ ) (Figure 19B) treated embryos, respectively. Cell cycle analysis was performed by analysis of approximately 8,000 events per condition.

Figures 20A-20C represent CAM tissue from CSAT (anti- $\beta_1$ ) treated embryos and Figures 20D-20F represent CAM tissue from LM609 (anti- $\alpha_v\beta_3$ ) treated embryos prepared as described in Example 12C. Figures 20A and 20D depict tissues stained with Apop Tag and visualized by fluorescence (FITC) superimposed on a D.I.C. image. Figures 20B and 20E depict the same tissues stained with mAb LM609 (anti- $\alpha_v\beta_3$ ) and visualized by fluorescence (rhodamine). Figures 20C and 20F represent merged images of the same tissues stained with both Apop Tag and LM609 where yellow staining represents colocalization. The bar represents 15 and 50  $\mu$ m in the left and right panels, respectively.

Detailed Description of the InventionA. Definitions

Amino Acid Residue: An amino acid formed upon chemical digestion (hydrolysis) of a polypeptide at its peptide linkages. The amino acid residues described herein are preferably in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the desired functional property is retained by the polypeptide. NH<sub>2</sub> refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxy terminus of a polypeptide. In keeping with standard polypeptide nomenclature (described in J. Biol. Chem., 243:3552-59 (1969) and adopted at 37 CFR §1.822(b)(2)), abbreviations for amino acid residues are shown in the following Table of Correspondence:

TABLE OF CORRESPONDENCE

	<u>SYMBOL</u>		<u>AMINO ACID</u>
	<u>1-Letter</u>	<u>3-Letter</u>	
	Y	Tyr	tyrosine
	G	Gly	glycine
25	F	Phe	phenylalanine
	M	Met	methionine
	A	Ala	alanine
	S	Ser	serine
	I	Ile	isoleucine
30	L	Leu	leucine
	T	Thr	threonine
	V	Val	valine
	P	Pro	proline
	K	Lys	lysine
35	H	His	histidine
	Q	Gln	glutamine
	E	Glu	glutamic acid
	Z	Glx	Glu and/or Gln

- 13 -

	W	Trp	tryptophan
	R	Arg	arginine
	D	Asp	aspartic acid
	N	Asn	asparagine
5	B	Asx	Asn and/or Asp
	C	Cys	cysteine
	X	Xaa	unknown/other

In addition the following have the meanings below:

	BOC	tert-butyloxycarbonyl
10	DCCI	dicylcohexylcarbodiimide
	DMF	dimethylformamide
	OMe	methoxy
	HOBT	1-hydroxybezotriazole

15           It should be noted that all amino acid residue sequences are represented herein by formulae whose left and right orientation is in the conventional direction of amino-terminus to carboxy-terminus. Furthermore, it should be noted that a dash at the  
20           beginning or end of an amino acid residue sequence indicates a peptide bond to a further sequence of one or more amino acid residues.

Polypeptide: refers to a linear series of amino acid residues connected to one another by  
25           peptide bonds between the alpha-amino group and carboxy group of contiguous amino acid residues.

Peptide: as used herein refers to a linear series of no more than about 50 amino acid residues connected one to the other as in a polypeptide.

30           Cyclic peptide: is derived from a corresponding linear peptide and; refers to a peptide in which no free N- or C-termini exist and; and of which the corresponding linear peptide's N-termini forms an amide bond to the C-terminal  
35           carboxylate of the said corresponding linear peptide.



- 14 -

Protein: refers to a linear series of greater than 50 amino acid residues connected one to the other as in a polypeptide.

5       Synthetic peptide: refers to a chemically produced chain of amino acid residues linked together by peptide bonds that is free of naturally occurring proteins and fragments thereof.

B.   General Considerations

10           The present invention relates generally to the discovery that angiogenesis is mediated by the specific vitronectin receptor  $\alpha_v\beta_3$ , and that inhibition of  $\alpha_v\beta_3$  function inhibits angiogenesis. This discovery is important because of the role  
15       that angiogenesis plays in a variety of disease processes. By inhibiting angiogenesis, one can intervene in the disease, ameliorate the symptoms, and in some cases cure the disease.

20           Where the growth of new blood vessels is the cause of, or contributes to, the pathology associated with a disease, inhibition of angiogenesis will reduce the deleterious effects of the disease. Examples include rheumatoid  
25       arthritis, diabetic retinopathy, inflammatory diseases, restenosis, and the like. Where the growth of new blood vessels is required to support growth of a deleterious tissue, inhibition of angiogenesis will reduce the blood supply to the tissue and thereby contribute to reduction in  
30       tissue mass based on blood supply requirements. Examples include growth of tumors where neovascularization is a continual requirement in order that the tumor grow beyond a few millimeters in thickness, and for the establishment of solid  
35       tumor metastases.

          The methods of the present invention are effective in part because the therapy is highly selective for angiogenesis and not other biological

- 15 -

processes. As shown in the Examples, only new vessel growth contains substantial  $\alpha_v\beta_3$ , and therefore the therapeutic methods do not adversely effect mature vessels. Furthermore,  $\alpha_v\beta_3$  is not widely distributed in normal tissues, but rather is found selectively on new vessels, thereby assuring that the therapy can be selectively targeted to new vessel growth.

The discovery that inhibition of  $\alpha_v\beta_3$  alone will effectively inhibit angiogenesis allows for the development of therapeutic compositions with potentially high specificity, and therefore relatively low toxicity. Thus although the invention discloses the use of peptide-based reagents which have the ability to inhibit one or more integrins, one can design other reagents which more selectively inhibit  $\alpha_v\beta_3$ , and therefore do not have the side effect of inhibiting other biological processes other than those mediated by  $\alpha_v\beta_3$ .

For example, as shown by the present teachings, it is possible to prepare monoclonal antibodies highly selective for immunoreaction with  $\alpha_v\beta_3$ , that are similarly selective for inhibition of  $\alpha_v\beta_3$  function. In addition, RGD-containing peptides can be designed to be selective for inhibition of  $\alpha_v\beta_3$ , as described further herein.

Prior to the discoveries of the present invention, it was not known that angiogenesis, and any of the processes dependent on angiogenesis, could be inhibited in vivo by the use of reagents that antagonize the biological function of  $\alpha_v\beta_3$ .

### C. Methods For Inhibition of Angiogenesis

The invention provides for a method for the inhibition of angiogenesis in a tissue, and thereby inhibiting events in the tissue which depend upon angiogenesis. Generally, the method comprises administering to the tissue a composition

- 16 -

comprising an angiogenesis-inhibiting amount of an  $\alpha_v\beta_3$  antagonist.

5 As described earlier, angiogenesis includes a variety of processes involving neovascularization of a tissue including "sprouting", vasculogenesis, or vessel enlargement, all of which angiogenesis processes are mediated by and dependent upon the expression of  $\alpha_v\beta_3$ . With the exception of traumatic wound healing, corpus leuteum formation and  
10 embryogenesis, it is believed that the majority of angiogenesis processes are associated with disease processes and therefore the use of the present therapeutic methods are selective for the disease and do not have deleterious side effects.

15 There are a variety of diseases in which angiogenesis is believed to be important, referred to as angiogenic diseases, including but not limited to, inflammatory disorders such as immune and non-immune inflammation, chronic articular  
20 rheumatism and psoriasis, disorders associated with inappropriate or inopportune invasion of vessels such as diabetic retinopathy, neovascular glaucoma, restenosis, capillary proliferation in atherosclerotic plaques and osteoporosis, and  
25 cancer associated disorders, such as solid tumors, solid tumor metastases, angiofibromas, retrolental fibroplasia, hemangiomas, Kaposi sarcoma and the like cancers which require neovascularization to support tumor growth.

30 Thus, methods which inhibit angiogenesis in a diseased tissue ameliorates symptoms of the disease and, depending upon the disease, can contribute to cure of the disease. In one embodiment, the invention contemplates inhibition of angiogenesis, per se, in a tissue. The extent of angiogenesis in  
35 a tissue, and therefore the extent of inhibition achieved by the present methods, can be evaluated by a variety of methods, such as are described in

- 17 -

the Examples for detecting  $\alpha_v\beta_3$ -immunopositive immature and nascent vessel structures by immunohistochemistry.

As described herein, any of a variety of tissues, or organs comprised of organized tissues, can support angiogenesis in disease conditions including skin, muscle, gut, connective tissue, joints, bones and the like tissue in which blood vessels can invade upon angiogenic stimuli.

Thus, in one related embodiment, a tissue to be treated is an inflamed tissue and the angiogenesis to be inhibited is inflamed tissue angiogenesis where there is neovascularization of inflamed tissue. In this class the method contemplates inhibition of angiogenesis in arthritic tissues, such as in a patient with chronic articular rheumatism, in immune or non-immune inflamed tissues, in psoriatic tissue and the like.

The patient treated in the present invention in its many embodiments is desirably a human patient, although it is to be understood that the principles of the invention indicate that the invention is effective with respect to all mammals, which are intended to be included in the term "patient". In this context, a mammal is understood to include any mammalian species in which treatment of diseases associated with angiogenesis is desirable, particularly agricultural and domestic mammalian species.

In another related embodiment, a tissue to be treated is a retinal tissue of a patient with diabetic retinopathy, macular degeneration or neovascular glaucoma and the angiogenesis to be inhibited is retinal tissue angiogenesis where there is neovascularization of retinal tissue.

In an additional related embodiment, a tissue to be treated is a tumor tissue of a patient with a

- 18 -

solid tumor, a metastases, a skin cancer, a breast cancer, a hemangioma or angiofibroma and the like cancer, and the angiogenesis to be inhibited is tumor tissue angiogenesis where there is

5 neovascularization of a tumor tissue. Typical solid tumor tissues treatable by the present methods include lung, pancreas, breast, colon, laryngeal, ovarian, and the like tissues. Exemplary tumor tissue angiogenesis, and inhibition

10 thereof, is described in the Examples.

Inhibition of tumor tissue angiogenesis is a particularly preferred embodiment because of the important role neovascularization plays in tumor growth. In the absence of neovascularization of

15 tumor tissue, the tumor tissue does not obtain the required nutrients, slows in growth, ceases additional growth, regresses and ultimately becomes necrotic resulting in killing of the tumor.

Stated in other words, the present invention provides for a method of inhibiting tumor

20 neovascularization by inhibiting tumor angiogenesis according to the present methods. Similarly, the invention provides a method of inhibiting tumor growth by practicing the angiogenesis-inhibiting

25 methods.

The methods are also particularly effective against the formation of metastases because (1) their formation requires vascularization of a primary tumor so that the metastatic cancer cells

30 can exit the primary tumor and (2) their establishment in a secondary site requires neovascularization to support growth of the metastases.

In a related embodiment, the invention

35 contemplates the practice of the method in conjunction with other therapies such as conventional chemotherapy directed against solid tumors and for control of establishment of

- 19 -

metastases. The administration of angiogenesis inhibitor is typically conducted during or after chemotherapy, although it is preferably to inhibit angiogenesis after a regimen of chemotherapy at times where the tumor tissue will be responding to the toxic assault by inducing angiogenesis to recover by the provision of a blood supply and nutrients to the tumor tissue. In addition, it is preferred to administer the angiogenesis inhibition methods after surgery where solid tumors have been removed as a prophylaxis against metastases.

Insofar as the present methods apply to inhibition of tumor neovascularization, the methods can also apply to inhibition of tumor tissue growth, to inhibition of tumor metastases formation, and to regression of established tumors. The Examples demonstrate regression of an established tumor following a single intravenous administration of an  $\alpha_v\beta_3$  antagonist of this invention.

Restenosis is a process of smooth muscle cell (SMC) migration and proliferation at the site of percutaneous transluminal coronary angioplasty which hampers the success of angioplasty. The migration and proliferation of SMC's during restenosis can be considered a process of angiogenesis which is inhibited by the present methods. Therefore, the invention also contemplates inhibition of restenosis by inhibiting angiogenesis according to the present methods in a patient following angioplasty procedures. For inhibition of restenosis, the  $\alpha_v\beta_3$  antagonist is typically administered after the angioplasty procedure for from about 2 to about 28 days, and more typically for about the first 14 days following the procedure.

The present method for inhibiting angiogenesis in a tissue, and therefore for also practicing the

- 20 -

methods for treatment of angiogenesis-related diseases, comprises contacting a tissue in which angiogenesis is occurring, or is at risk for occurring, with a composition comprising a therapeutically effective amount of an  $\alpha_v\beta_3$  antagonist capable of inhibiting  $\alpha_v\beta_3$  binding to its natural ligand. Thus the method comprises administering to a patient a therapeutically effective amount of a physiologically tolerable composition containing an  $\alpha_v\beta_3$  antagonist of the invention.

The dosage ranges for the administration of the  $\alpha_v\beta_3$  antagonist depend upon the form of the antagonist, and its potency, as described further herein, and are amounts large enough to produce the desired effect in which angiogenesis and the disease symptoms mediated by angiogenesis are ameliorated. The dosage should not be so large as to cause adverse side effects, such as hyperviscosity syndromes, pulmonary edema, congestive heart failure, and the like. Generally, the dosage will vary with the age, condition, sex and extent of the disease in the patient and can be determined by one of skill in the art. The dosage can also be adjusted by the individual physician in the event of any complication.

A therapeutically effective amount is an amount of  $\alpha_v\beta_3$  antagonist sufficient to produce a measurable inhibition of angiogenesis in the tissue being treated, ie., an angiogenesis-inhibiting amount. Inhibition of angiogenesis can be measured in situ by immunohistochemistry, as described herein, or by other methods known to one skilled in the art.

Insofar as an  $\alpha_v\beta_3$  antagonist can take the form of a  $\alpha_v\beta_3$  mimetic, an RGD-containing peptide, an anti- $\alpha_v\beta_3$  monoclonal antibody, or fragment thereof, it is to be appreciated that the potency, and

- 21 -

therefore an expression of a "therapeutically effective" amount can vary. However, as shown by the present assay methods, one skilled in the art can readily assess the potency of a candidate  $\alpha_v\beta_3$  antagonist of this invention.

Potency of an  $\alpha_v\beta_3$  antagonist can be measured by a variety of means including inhibition of angiogenesis in the CAM assay, in the in vivo rabbit eye assay, in the in vivo chimeric mouse:human assay, and by measuring inhibition of binding of natural ligand to  $\alpha_v\beta_3$ , all as described herein, and the like assays.

A preferred  $\alpha_v\beta_3$  antagonist has the ability to substantially inhibit binding of a natural ligand such as fibrinogen or vitronectin to  $\alpha_v\beta_3$  in solution at antagonist concentrations of less than 0.5 micromolar ( $\mu\text{M}$ ), preferably less than 0.1  $\mu\text{M}$ , and more preferably less than 0.05  $\mu\text{M}$ . By "substantially" is meant that at least a 50 percent reduction in binding of fibrinogen is observed by inhibition in the presence of the  $\alpha_v\beta_3$  antagonist, and at 50% inhibition is referred to herein as an  $\text{IC}_{50}$  value.

A more preferred  $\alpha_v\beta_3$  antagonist exhibits selectivity for  $\alpha_v\beta_3$  over other integrins. Thus, a preferred  $\alpha_v\beta_3$  antagonist substantially inhibits fibrinogen binding to  $\alpha_v\beta_3$  but does not substantially inhibit binding of fibrinogen to another integrin, such as  $\alpha_v\beta_1$ ,  $\alpha_v\beta_5$  or  $\alpha_{\text{IIb}}\beta_3$ . Particularly preferred is an  $\alpha_v\beta_3$  antagonist that exhibits a 10-fold to 100-fold lower  $\text{IC}_{50}$  activity at inhibiting fibrinogen binding to  $\alpha_v\beta_3$  compared to the  $\text{IC}_{50}$  activity at inhibiting fibrinogen binding to another integrin. Exemplary assays for measuring  $\text{IC}_{50}$  activity at inhibiting fibrinogen binding to an integrin are described in the Examples.



- 22 -

A therapeutically effective amount of an  $\alpha_v\beta_3$  antagonist of this invention in the form of a monoclonal antibody is typically an amount such that when administered in a physiologically tolerable composition is sufficient to achieve a plasma concentration of from about 0.01 microgram (ug) per milliliter (ml) to about 100 ug/ml, preferably from about 1 ug/ml to about 5 ug/ml, and usually about 5 ug/ml. Stated differently, the dosage can vary from about 0.1 mg/kg to about 300 mg/kg, preferably from about 0.2 mg/kg to about 200 mg/kg, most preferably from about 0.5 mg/kg to about 20 mg/kg, in one or more dose administrations daily, for one or several days.

Where the antagonist is in the form of a fragment of a monoclonal antibody, the amount can readily be adjusted based on the mass of the fragment relative to the mass of the whole antibody. A preferred plasma concentration in molarity is from about 2 micromolar ( $\mu$ M) to about 5 millimolar (mM) and preferably about 100  $\mu$ M to 1 mM antibody antagonist.

A therapeutically effective amount of an  $\alpha_v\beta_3$  antagonist of this invention in the form of a polypeptide, or other similarly-sized small molecule  $\alpha_v\beta_3$  mimetic, is typically an amount of polypeptide such that when administered in a physiologically tolerable composition is sufficient to achieve a plasma concentration of from about 0.1 microgram (ug) per milliliter (ml) to about 200 ug/ml, preferably from about 1 ug/ml to about 150 ug/ml. Based on a polypeptide having a mass of about 500 grams per mole, the preferred plasma concentration in molarity is from about 2 micromolar ( $\mu$ M) to about 5 millimolar (mM) and preferably about 100  $\mu$ M to 1 mM polypeptide antagonist. Stated differently, the dosage per body weight can vary from about 0.1 mg/kg to about

- 23 -

300 mg/kg, and preferably from about 0.2 mg/kg to about 200 mg/kg, in one or more dose administrations daily, for one or several days.

5 The monoclonal antibodies or polypeptides of the invention can be administered parenterally by injection or by gradual infusion over time. Although the tissue to be treated can typically be accessed in the body by systemic administration and therefore most often treated by intravenous  
10 administration of therapeutic compositions, other tissues and delivery means are contemplated where there is a likelihood that the tissue targeted contains the target molecule. Thus, monoclonal antibodies or polypeptides of the invention can be  
15 administered intravenously, intraperitoneally, intramuscularly, subcutaneously, intracavity, transdermally, and can be delivered by peristaltic means.

The therapeutic compositions containing a  
20 monoclonal antibody or a polypeptide of this invention are conventionally administered intravenously, as by injection of a unit dose, for example. The term "unit dose" when used in reference to a therapeutic composition of the  
25 present invention refers to physically discrete units suitable as unitary dosage for the subject, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required  
30 diluent; i.e., carrier, or vehicle.

In one preferred embodiment as shown in the Examples, the  $\alpha_v\beta_3$  antagonist is administered in a single dosage intravenously.

35 The compositions are administered in a manner compatible with the dosage formulation, and in a therapeutically effective amount. The quantity to be administered and timing depends on the subject to be treated, capacity of the subject's system to

- 24 -

utilize the active ingredient, and degree of therapeutic effect desired. Precise amounts of active ingredient required to be administered depend on the judgement of the practitioner and are peculiar to each individual. However, suitable dosage ranges for systemic application are disclosed herein and depend on the route of administration. Suitable regimes for administration are also variable, but are typified by an initial administration followed by repeated doses at one or more hour intervals by a subsequent injection or other administration. Alternatively, continuous intravenous infusion sufficient to maintain concentrations in the blood in the ranges specified for in vivo therapies are contemplated.

As demonstrated by the present Examples, inhibition of angiogenesis and tumor regression occurs as early as 7 days after the initial contacting with antagonist. Additional or prolonged exposure to antagonist is preferable for 7 days to 6 weeks, preferably about 14 to 28 days.

In a related embodiment, the Examples demonstrate the relationship between inhibition of  $\alpha_v\beta_3$  and induction of apoptosis in the neovasculature cells bearing  $\alpha_v\beta_3$ . Thus, the invention also contemplates methods for inhibition of apoptosis in neovasculature of a tissue. The method is practiced substantially as described herein for inhibition of angiogenesis in all tissues and conditions described therefor. The only noticeable difference is one of timing of effect, which is that apoptosis is manifest quickly, typically about 48 hours after contacting antagonist, whereas inhibition of angiogenesis and tumor regression is manifest more slowly, as described herein. This difference affects the therapeutic regimen in terms of time of administration, and effect desired. Typically,

- 25 -

administration for apoptosis of neovasculature can be for 24 hours to about 4 weeks, although 48 hours to 7 days is preferred.

5           D.   Therapeutic Compositions

          The present invention contemplates therapeutic compositions useful for practicing the therapeutic methods described herein. Therapeutic compositions of the present invention contain a  
10           physiologically tolerable carrier together with an  $\alpha_v\beta_3$  antagonist as described herein, dissolved or dispersed therein as an active ingredient. In a preferred embodiment, the therapeutic  $\alpha_v\beta_3$  antagonist composition is not immunogenic when  
15           administered to a mammal or human patient for therapeutic purposes.

          As used herein, the terms "pharmaceutically acceptable", "physiologically tolerable" and grammatical variations thereof, as they refer to  
20           compositions, carriers, diluents and reagents, are used interchangeably and represent that the materials are capable of administration to or upon a mammal without the production of undesirable physiological effects such as nausea, dizziness,  
25           gastric upset and the like.

          The preparation of a pharmacological composition that contains active ingredients dissolved or dispersed therein is well understood in the art and need not be limited based on  
30           formulation. Typically such compositions are prepared as injectables either as liquid solutions or suspensions, however, solid forms suitable for solution, or suspensions, in liquid prior to use can also be prepared. The preparation can also be  
35           emulsified.

---

          The active ingredient can be mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient and in

amounts suitable for use in the therapeutic methods described herein. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol or the like and combinations thereof. In addition, if desired, the composition can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like which enhance the effectiveness of the active ingredient.

The therapeutic composition of the present invention can include pharmaceutically acceptable salts of the components therein. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide) that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, tartaric, mandelic and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine and the like.

Particularly preferred are the salts of TFA and HCl, when used in the preparation of cyclic polypeptide  $\alpha, \beta$ , antagonists. Representative salts of peptides are described in the Examples.

Physiologically tolerable carriers are well known in the art. Exemplary of liquid carriers are sterile aqueous solutions that contain no materials in addition to the active ingredients and water, or contain a buffer such as sodium phosphate at physiological pH value, physiological saline or both, such as phosphate-buffered saline. Still further, aqueous carriers can contain more than one buffer salt, as well as salts such as sodium and

- 27 -

potassium chlorides, dextrose, polyethylene glycol and other solutes.

Liquid compositions can also contain liquid phases in addition to and to the exclusion of water. Exemplary of such additional liquid phases are glycerin, vegetable oils such as cottonseed oil, and water-oil emulsions.

A therapeutic composition contains an angiogenesis-inhibiting amount of an  $\alpha_v\beta_3$  antagonist of the present invention, typically formulated to contain an amount of at least 0.1 weight percent of antagonist per weight of total therapeutic composition. A weight percent is a ratio by weight of inhibitor to total composition. Thus, for example, 0.1 weight percent is 0.1 grams of inhibitor per 100 grams of total composition.

E. Antagonists of Integrin  $\alpha_v\beta_3$

$\alpha_v\beta_3$  antagonists are used in the present methods for inhibiting angiogenesis in tissues, and can take a variety of forms that include compounds which interact with  $\alpha_v\beta_3$  in a manner such that functional interactions with natural  $\alpha_v\beta_3$  ligands are interfered. Exemplary antagonists include analogs of  $\alpha_v\beta_3$  derived from the ligand binding site on  $\alpha_v\beta_3$ , mimetics of either  $\alpha_v\beta_3$  or a natural ligand of  $\alpha_v\beta_3$  that mimic the structural region involved in  $\alpha_v\beta_3$ -ligand binding interactions, polypeptides having a sequence corresponding to a functional binding domain of the natural ligand specific for  $\alpha_v\beta_3$ , particularly corresponding to the RGD-containing domain of a natural ligand of  $\alpha_v\beta_3$ , and antibodies which immunoreact with either  $\alpha_v\beta_3$  or the natural ligand, all of which exhibit antagonist activity as defined herein.

### 1. Polypeptides

In one embodiment, the invention contemplates  $\alpha_v\beta_3$  antagonists in the form of polypeptides. A polypeptide (peptide)  $\alpha_v\beta_3$  antagonist can have the sequence characteristics of either the natural ligand of  $\alpha_v\beta_3$  or  $\alpha_v\beta_3$  itself at the region involved in  $\alpha_v\beta_3$ -ligand interaction and exhibits  $\alpha_v\beta_3$  antagonist activity as described herein. A preferred  $\alpha_v\beta_3$  antagonist peptide contains the RGD tripeptide and corresponds in sequence to the natural ligand in the RGD-containing region.

Preferred RGD-containing polypeptides have a sequence corresponding to the amino acid residue sequence of the RGD-containing region of a natural ligand of  $\alpha_v\beta_3$ , such as fibrinogen, vitronectin, von Willebrand factor, laminin, thrombospondin, and the like ligands. The sequence of these  $\alpha_v\beta_3$  ligands are well known. Thus, an  $\alpha_v\beta_3$  antagonist peptide can be derived from any of the natural ligands, although fibrinogen and vitronectin are preferred.

A particularly preferred  $\alpha_v\beta_3$  antagonist peptide preferentially inhibits  $\alpha_v\beta_3$  binding to its natural ligand(s) when compared to other integrins, as described earlier. These  $\alpha_v\beta_3$ -specific peptides are particularly preferred at least because the specificity for  $\alpha_v\beta_3$  reduces the incidence of undesirable side effects such as inhibition of other integrins. The identification of preferred  $\alpha_v\beta_3$  antagonist peptides having selectivity for  $\alpha_v\beta_3$  can readily be identified in a typical inhibition of binding assay, such as the ELISA assay described in the Examples.

In one embodiment, a polypeptide of the present invention comprises no more than about 100 amino acid residues, preferably no more than about 60 residues, more preferably no more than about 30 residues. Peptides can be linear or cyclic,

- 29 -

although particularly preferred peptides are cyclic.

Preferred cyclic and linear peptides and their designations are shown in Table 1 in the Examples.

5 It should be understood that a subject polypeptide need not be identical to the amino acid residue sequence of a  $\alpha_v\beta_3$  natural ligand, so long as it includes the required sequence and is able to function as an  $\alpha_v\beta_3$  antagonist in an assay such as those described herein.

10 A subject polypeptide includes any analog, fragment or chemical derivative of a polypeptide whose amino acid residue sequence is shown herein so long as the polypeptide is an  $\alpha_v\beta_3$  antagonist. Therefore, a present polypeptide can be subject to various changes, substitutions, insertions, and deletions where such changes provide for certain advantages in its use. In this regard,  $\alpha_v\beta_3$  antagonist polypeptide of this invention

15 corresponds to, rather than is identical to, the sequence of a recited peptide where one or more changes are made and it retains the ability to function as an  $\alpha_v\beta_3$  antagonist in one or more of the assays as defined herein.

20 Thus, a polypeptide can be in any of a variety of forms of peptide derivatives, that include amides, conjugates with proteins, cyclized peptides, polymerized peptides, analogs, fragments, chemically modified peptides, and the like derivatives.

25 The term "analog" includes any polypeptide having an amino acid residue sequence substantially identical to a sequence specifically shown herein in which one or more residues have been conservatively substituted with a functionally

30 similar residue and which displays the  $\alpha_v\beta_3$  antagonist activity as described herein. Examples of conservative substitutions include the



- 30 -

substitution of one non-polar (hydrophobic) residue such as isoleucine, valine, leucine or methionine for another, the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, between glycine and serine, the substitution of one basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue, such as aspartic acid or glutamic acid for another.

The phrase "conservative substitution" also includes the use of a chemically derivatized residue in place of a non-derivatized residue provided that such polypeptide displays the requisite inhibition activity.

"Chemical derivative" refers to a subject polypeptide having one or more residues chemically derivatized by reaction of a functional side group. Such derivatized molecules include for example, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene sulfonyl groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloroacetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and ethyl esters or other types of esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine may be derivatized to form N-im-benzylhistidine. Also included as chemical derivatives are those peptides which contain one or more naturally occurring amino acid derivatives of the twenty standard amino acids. For examples: 4-hydroxyproline may be substituted for proline; 5-hydroxylysine may be substituted for lysine; 3-methylhistidine may be substituted for histidine; homoserine may be substituted for serine; and ornithine may be substituted for lysine.

- 31 -

Polypeptides of the present invention also include any polypeptide having one or more additions and/or deletions or residues relative to the sequence of a polypeptide whose sequence is shown herein, so long as the requisite activity is maintained.

The term "fragment" refers to any subject polypeptide having an amino acid residue sequence shorter than that of a polypeptide whose amino acid residue sequence is shown herein.

When a polypeptide of the present invention has a sequence that is not identical to the sequence of an  $\alpha, \beta$ , natural ligand, it is typically because one or more conservative or non-conservative substitutions have been made, usually no more than about 30 number percent, and preferably no more than 10 number percent of the amino acid residues are substituted. Additional residues may also be added at either terminus of a polypeptide for the purpose of providing a "linker" by which the polypeptides of this invention can be conveniently affixed to a label or solid matrix, or carrier.

Labels, solid matrices and carriers that can be used with the polypeptides of this invention are described hereinbelow.

Amino acid residue linkers are usually at least one residue and can be 40 or more residues, more often 1 to 10 residues, but do not form  $\alpha, \beta$ , ligand epitopes. Typical amino acid residues used for linking are tyrosine, cysteine, lysine, glutamic and aspartic acid, or the like. In addition, a subject polypeptide can differ, unless otherwise specified, from the natural sequence of an  $\alpha, \beta$ , ligand by the sequence being modified by terminal-NH<sub>2</sub> acylation, e.g., acetylation, or thioglycolic acid amidation, by terminal-carboxylamidation, e.g., with ammonia, methylamine, and the like terminal modifications. Terminal

- 32 -

modifications are useful, as is well known, to reduce susceptibility by proteinase digestion, and therefore serve to prolong half life of the polypeptides in solutions, particularly biological fluids where proteases may be present. In this regard, polypeptide cyclization is also a useful terminal modification, and is particularly preferred also because of the stable structures formed by cyclization and in view of the biological activities observed for such cyclic peptides as described herein.

Any peptide of the present invention may be used in the form of a pharmaceutically acceptable salt. Suitable acids which are capable of forming salts with the peptides of the present invention include inorganic acids such as trifluoroacetic acid (TFA) hydrochloric acid (HCl), hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, phosphoric acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, malonic acid, succinic acid, maleic acid, fumaric acid, anthranilic acid, cinnamic acid, naphthalene sulfonic acid, sulfanilic acid or the like. HCl and TFA salts are particularly preferred.

Suitable bases capable of forming salts with the peptides of the present invention include inorganic bases such as sodium hydroxide, ammonium hydroxide, potassium hydroxide and the like; and organic bases such as mono-, di- and tri-alkyl and aryl amines (e.g. triethylamine, diisopropyl amine, methyl amine, dimethyl amine and the like) and optionally substituted ethanolamines (e.g. ethanolamine, diethanolamine and the like).

A peptide of the present invention also referred to herein as a subject polypeptide, can be synthesized by any of the techniques that are known to those skilled in the polypeptide art, including

- 33 -

recombinant DNA techniques. Synthetic chemistry techniques, such as a solid-phase Merrifield-type synthesis, are preferred for reasons of purity, antigenic specificity, freedom from undesired side products, ease of production and the like. An excellent summary of the many techniques available can be found in Steward et al., "Solid Phase Peptide Synthesis", W.H. Freeman Co., San Francisco, 1969; Bodanszky, et al., "Peptide Synthesis", John Wiley & Sons, Second Edition, 1976; J. Meienhofer, "Hormonal Proteins and Peptides", Vol. 2, p. 46, Academic Press (New York), 1983; Merrifield, Adv. Enzymol., 32:221-96, 1969; Fields et al., Int. J. Peptide Protein Res., 35:161-214, 1990; and United States Patent No. 4,244,946 for solid phase peptide synthesis, and Schroder et al., "The Peptides", Vol. 1, Academic Press (New York), 1965 for classical solution synthesis, each of which is incorporated herein by reference. Appropriate protective groups usable in such synthesis are described in the above texts and in J.F.W. McOmie, "Protective Groups in Organic Chemistry", Plenum Press, New York, 1973, which is incorporated herein by reference.

In general, the solid-phase synthesis methods contemplated comprise the sequential addition of one or more amino acid residues or suitably protected amino acid residues to a growing peptide chain. Normally, either the amino or carboxyl group of the first amino acid residue is protected by a suitable, selectively removable protecting group. A different, selectively removable protecting group is utilized for amino acids containing a reactive side group such as lysine.

Using a solid phase synthesis as exemplary, the protected or derivatized amino acid is attached to an inert solid support through its unprotected carboxyl or amino group. The protecting group of

- 34 -

the amino or carboxyl group is then selectively removed and the next amino acid in the sequence having the complimentary (amino or carboxyl) group suitably protected is admixed and reacted under conditions suitable for forming the amide linkage with the residue already attached to the solid support. The protecting group of the amino or carboxyl group is then removed from this newly added amino acid residue, and the next amino acid (suitably protected) is then added, and so forth. After all the desired amino acids have been linked in the proper sequence, any remaining terminal and side group protecting groups (and solid support) are removed sequentially or concurrently, to afford the final linear polypeptide.

The resultant linear polypeptides prepared for example as described above may be reacted to form their corresponding cyclic peptides. An exemplary method for cyclizing peptides is described by Zimmer et al., Peptides 1992, pp. 393-394, ESCOM Science Publishers, B.V., 1993. Typically, tertbutoxycarbonyl protected peptide methyl ester is dissolved in methanol and sodium hydroxide solution are added and the admixture is reacted at 20°C (20C) to hydrolytically remove the methyl ester protecting group. After evaporating the solvent, the tertbutoxycarbonyl protected peptide is extracted with ethyl acetate from acidified aqueous solvent. The tertbutoxycarbonyl protecting group is then removed under mildly acidic conditions in dioxane cosolvent. The unprotected linear peptide with free amino and carboxy termini so obtained is converted to its corresponding cyclic peptide by reacting a dilute solution of the linear peptide, in a mixture of dichloromethane and dimethylformamide, with dicyclohexylcarbodiimide in the presence of 1-hydroxybenzotriazole and N-

- 35 -

methyilmorpholine. The resultant cyclic peptide is then purified by chromatography.

A particularly preferred cyclic peptide synthesis method is described by Gurrath et al.,  
5 Eur. J. Biochem., 210:911-921 (1992), and described in the Examples. Particularly preferred peptides for use in the present methods are c-(GrGDFV) (SEQ ID NO 4), c-(RGDFV) (SEQ ID NO 5), c-(RADfV) (SEQ ID NO 6), c-(RGDFv) (SEQ ID NO 7) and linear  
10 peptide YTAECKPQVTRGDVF (SEQ ID NO 8), where "c-" indicates a cyclic peptide, the upper case letters are single letter code for an L-amino acid and the lower case letters are single letter code for D-amino acid. The amino acid residues sequence of  
15 these peptides are also shown in SEQ ID NOS 4, 5, 6, 7 and 8, respectively.

## 2. Monoclonal Antibodies

The present invention describes, in  
20 one embodiment,  $\alpha_v\beta_3$  antagonists in the form of monoclonal antibodies which immunoreact with  $\alpha_v\beta_3$  and inhibit  $\alpha_v\beta_3$  binding to its natural ligand as described herein. The invention also describes cell lines which produce the antibodies, methods  
25 for producing the cell lines, and methods for producing the monoclonal antibodies.

A monoclonal antibody of this invention comprises antibody molecules that 1) immunoreact with isolated  $\alpha_v\beta_3$ , and 2) inhibit fibrinogen  
30 binding to  $\alpha_v\beta_3$ . Preferred monoclonal antibodies which preferentially bind to  $\alpha_v\beta_3$  include a monoclonal antibody having the immunoreaction characteristics of mAb LM609, secreted by hybridoma cell line ATCC HB 9537. The hybridoma cell line  
35 ATCC HB 9537 was deposited pursuant to Budapest Treaty requirements with the American Type Culture Collection (ATCC), 1301 Parklawn Drive, Rockville, MD, USA, on September 15, 1987.

- 36 -

The term "antibody or antibody molecule" in the various grammatical forms is used herein as a collective noun that refers to a population of immunoglobulin molecules and/or immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antibody combining site or paratope.

An "antibody combining site" is that structural portion of an antibody molecule comprised of heavy and light chain variable and hypervariable regions that specifically binds antigen.

Exemplary antibodies for use in the present invention are intact immunoglobulin molecules, substantially intact immunoglobulin molecules and those portions of an immunoglobulin molecule that contain the paratope, including those portions known in the art as Fab, Fab', F(ab')<sub>2</sub>, and F(v), and also referred to as antibody fragments.

In another preferred embodiment, the invention contemplates a truncated immunoglobulin molecule comprising a Fab fragment derived from a monoclonal antibody of this invention. The Fab fragment, lacking Fc receptor, is soluble, and affords therapeutic advantages in serum half life, and diagnostic advantages in modes of using the soluble Fab fragment. The preparation of a soluble Fab fragment is generally known in the immunological arts and can be accomplished by a variety of methods.

For example, Fab and F(ab')<sub>2</sub> portions (fragments) of antibodies are prepared by the proteolytic reaction of papain and pepsin, respectively, on substantially intact antibodies by methods that are well known. See for example, U.S. Patent No. 4,342,566 to Theofilopolous and Dixon. Fab' antibody portions are also well known and are produced from F(ab')<sub>2</sub> portions followed by reduction

- 37 -

of the disulfide bonds linking the two heavy chain portions as with mercaptoethanol, and followed by alkylation of the resulting protein mercaptan with a reagent such as iodoacetamide. An antibody containing intact immunoglobulin molecules are preferred, and are utilized as illustrative herein.

The phrase "monoclonal antibody" in its various grammatical forms refers to a population of antibody molecules that contain only one species of antibody combining site capable of immunoreacting with a particular epitope. A monoclonal antibody thus typically displays a single binding affinity for any epitope with which it immunoreacts. A monoclonal antibody may therefore contain an antibody molecule having a plurality of antibody combining sites, each immunospecific for a different epitope, e.g., a bispecific monoclonal antibody.

A monoclonal antibody is typically composed of antibodies produced by clones of a single cell called a hybridoma that secretes (produces) only one kind of antibody molecule. The hybridoma cell is formed by fusing an antibody-producing cell and a myeloma or other self-perpetuating cell line. The preparation of such antibodies was first described by Kohler and Milstein, Nature 256:495-497 (1975), which description is incorporated by reference. Additional methods are described by Zola, Monoclonal Antibodies: A Manual of Techniques, CRC Press, Inc. (1987). The hybridoma supernates so prepared can be screened for the presence of antibody molecules that immunoreact with  $\alpha_v\beta_3$ , and for inhibition of  $\alpha_v\beta_3$  binding to natural ligands.

Briefly, to form the hybridoma from which the monoclonal antibody composition is produced, a myeloma or other self-perpetuating cell line is fused with lymphocytes obtained from the spleen of



- 38 -

a mammal hyperimmunized with a source of  $\alpha_v\beta_3$ , such as  $\alpha_v\beta_3$  isolated from M21 human melanoma cells as described by Cheresch et al., J. Biol. Chem., 262:17703-17711 (1987).

5           It is preferred that the myeloma cell line used to prepare a hybridoma be from the same species as the lymphocytes. Typically, a mouse of the strain 129 GLX<sup>+</sup> is the preferred mammal. Suitable mouse myelomas for use in the present  
10          invention include the hypoxanthine-aminopterin-thymidine-sensitive (HAT) cell lines P3X63-Ag8.653, and Sp2/0-Ag14 that are available from the American Type Culture Collection, Rockville, MD, under the designations CRL 1580 and CRL 1581, respectively.

15           Splenocytes are typically fused with myeloma cells using polyethylene glycol (PEG) 1500. Fused hybrids are selected by their sensitivity to HAT. Hybridomas producing a monoclonal antibody of this invention are identified using the enzyme linked  
20          immunosorbent assay (ELISA) described in the Examples.

          A monoclonal antibody of the present invention can also be produced by initiating a monoclonal hybridoma culture comprising a nutrient medium  
25          containing a hybridoma that secretes antibody molecules of the appropriate specificity. The culture is maintained under conditions and for a time period sufficient for the hybridoma to secrete the antibody molecules into the medium. The  
30          antibody-containing medium is then collected. The antibody molecules can then be further isolated by well known techniques.

          Media useful for the preparation of these compositions are both well known in the art and  
35          commercially available and include synthetic culture media, inbred mice and the like. An exemplary synthetic medium is Dulbecco's minimal essential medium (DMEM; Dulbecco et al., Virology.

- 39 -

8:396, 1959) supplemented with 4.5 gm/l glucose, 20 mM glutamine, and 20% fetal calf serum. An exemplary inbred mouse strain is the Balb/c.

Other methods of producing a monoclonal antibody, a hybridoma cell, or a hybridoma cell culture are also well known. See, for example, the method of isolating monoclonal antibodies from an immunological repertoire as described by Sastry, et al., Proc. Natl. Acad. Sci. USA, 86:5728-5732 (1989); and Huse et al., Science, 246:1275-1281 (1989).

Also contemplated by this invention is the hybridoma cell, and cultures containing a hybridoma cell that produce a monoclonal antibody of this invention. Particularly preferred is the hybridoma cell line that secretes monoclonal antibody mAb LM609 designated ATCC HB 9537. mAb LM609 was prepared as described by Cheresh et al., J. Biol. Chem., 262:17703-17711 (1987), and its preparation is also described in the Examples.

The invention contemplates, in one embodiment, a monoclonal antibody that has the immunoreaction characteristics of mAb LM609.

It is also possible to determine, without undue experimentation, if a monoclonal antibody has the same (i.e., equivalent) specificity (immunoreaction characteristics) as a monoclonal antibody of this invention by ascertaining whether the former prevents the latter from binding to a preselected target molecule. If the monoclonal antibody being tested competes with the monoclonal antibody of the invention, as shown by a decrease in binding by the monoclonal antibody of the invention in standard competition assays for binding to the target molecule when present in the solid phase, then it is likely that the two monoclonal antibodies bind to the same, or a closely related, epitope.

- 40 -

Still another way to determine whether a monoclonal antibody has the specificity of a monoclonal antibody of the invention is to pre-incubate the monoclonal antibody of the invention with the target molecule with which it is normally reactive, and then add the monoclonal antibody being tested to determine if the monoclonal antibody being tested is inhibited in its ability to bind the target molecule. If the monoclonal antibody being tested is inhibited then, in all likelihood, it has the same, or functionally equivalent, epitopic specificity as the monoclonal antibody of the invention.

An additional way to determine whether a monoclonal antibody has the specificity of a monoclonal antibody of the invention is to determine the amino acid residue sequence of the CDR regions of the antibodies in question. Antibody molecules having identical, or functionally equivalent, amino acid residue sequences in their CDR regions have the same binding specificity. Methods for sequencing polypeptides is well known in the art.

The immunospecificity of an antibody, its target molecule binding capacity, and the attendant affinity the antibody exhibits for the epitope, are defined by the epitope with which the antibody immunoreacts. The epitope specificity is defined at least in part by the amino acid residue sequence of the variable region of the heavy chain of the immunoglobulin the antibody, and in part by the light chain variable region amino acid residue sequence.

Use of the term "having the binding specificity of" indicates that equivalent monoclonal antibodies exhibit the same or similar immunoreaction (binding) characteristics and

- 41 -

compete for binding to a preselected target molecule.

Humanized monoclonal antibodies offer particular advantages over murine monoclonal antibodies, particularly insofar as they can be used therapeutically in humans. Specifically, human antibodies are not cleared from the circulation as rapidly as "foreign" antigens, and do not activate the immune system in the same manner as foreign antigens and foreign antibodies. Methods of preparing "humanized" antibodies are generally well known in the art, and can readily be applied to the antibodies of the present invention.

Thus, the invention contemplates, in one embodiment, a monoclonal antibody of this invention that is humanized by grafting to introduce components of the human immune system without substantially interfering with the ability of the antibody to bind antigen.

### 3. $\alpha_v\beta_3$ -Specific Mimetics

The present invention demonstrates that  $\alpha_v\beta_3$  antagonists generally can be used in the present invention, which antagonists can include polypeptides, antibodies and other molecules, designated "mimetics", which have the capacity to interfere with  $\alpha_v\beta_3$  function. Particularly preferred are antagonists which specifically interfere with  $\alpha_v\beta_3$  function, and do not interfere with function of other integrins.

In this context it is appreciated that a variety of reagents may be suitable for use in the present methods, so long as these reagents possess the requisite biological activity. These reagents are generically referred to as mimetics because they possess the ability to "mimic" a binding domain on either  $\alpha_v\beta_3$ , or the  $\alpha_v\beta_3$  ligand involved in the functional interaction of the receptor and ligand,

- 42 -

and thereby interfere with (i.e., inhibit) normal function.

An  $\alpha_v\beta_3$  mimetic is any molecule, other than an antibody or ligand-derived peptide, which exhibits the above-described properties. It can be a synthetic analog of a peptide, a compound which is shaped like the binding pocket of the above-described binding domain, or other molecule.

The design of an  $\alpha_v\beta_3$  mimetic can be conducted by any of a variety of structural analysis methods for drug-design known in the art, including molecular modelling, two-dimensional nuclear magnetic resonance (2-D NMR) analysis, x-ray crystallography, random screening of peptide, peptide analog or other chemical polymer libraries, and the like drug design methodologies.

In view of the broad structural evidence presented in the present specification which shows that an  $\alpha_v\beta_3$  antagonist can be a small polypeptide or an monoclonal antibody, two diversely different chemical structures which share the functional property of selective inhibition of  $\alpha_v\beta_3$ , the structure of a subject  $\alpha_v\beta_3$  antagonist useful in the present methods need not be so limited, but includes any  $\alpha_v\beta_3$  mimetic, as defined herein.

F. Methods For Identifying Antagonists of  $\alpha_v\beta_3$

The invention also described assay methods for identifying candidate  $\alpha_v\beta_3$  antagonists for use according to the present methods. In these assay methods candidate molecules are evaluated for their potency in inhibiting  $\alpha_v\beta_3$  binding to natural ligands, and furthermore are evaluated for their potency in inhibiting angiogenesis in a tissue.

The first assay measures inhibition of direct binding of natural ligand to  $\alpha_v\beta_3$ , and a preferred embodiment is described in detail in the Examples.

- 43 -

The assay typically measures the degree of inhibition of binding of a natural ligand, such as fibrinogen, to isolated  $\alpha_v\beta_3$  in the solid phase by ELISA.

5           The assay can also be used to identify compounds which exhibit specificity for  $\alpha_v\beta_3$  and do not inhibit natural ligands from binding other integrins. The specificity assay is conducted by running parallel ELISA assays where both  $\alpha_v\beta_3$  and  
10           other integrins are screened concurrently in separate assay chambers for their respective abilities to bind a natural ligand and for the candidate compound to inhibit the respective  
15           abilities of the integrins to bind a preselected ligand. Preferred screening assay formats are described in the Examples.

          The second assay measures angiogenesis in the chick chorioallantoic membrane (CAM) and is referred to as the CAM assay. The CAM assay has  
20           been described in detail by others, and further has been used to measure both angiogenesis and neovascularization of tumor tissues. See Ausprunk et al., Am. J. Pathol., 79:597-618 (1975) and  
          Ossonski et al., Cancer Res., 40:2300-2309 (1980).

25           The CAM assay is a well recognized assay model for in vivo angiogenesis because neovascularization of whole tissue is occurring, and actual chick embryo blood vessels are growing into the CAM or into the tissue grown on the CAM.

30           As demonstrated herein, the CAM assay illustrates inhibition of neovascularization based on both the amount and extent of new vessel growth. Furthermore, it is easy to monitor the growth of any tissue transplanted upon the CAM, such as a  
35           tumor tissue. Finally, the assay is particularly useful because there is an internal control for toxicity in the assay system. The chick embryo is

- 44 -

exposed to any test reagent, and therefore the health of the embryo is an indication of toxicity.

5 The third assay measures angiogenesis is the in vivo rabbit eye model and is referred to as the rabbit eye assay. The rabbit eye assay has been described in detail by others, and further has been used to measure both angiogenesis and neovascularization in the presence of angiogenic inhibitors such as thalidomide. See D'Amato, et al., Proc. Natl. Acad. Sci., 91:4082-4085 (1994).

10 The rabbit eye assay is a well recognized assay model for in vivo angiogenesis because the neovascularization process, exemplified by rabbit blood vessels growing from the rim of the cornea into the cornea, is easily visualized through the naturally transparent cornea of the eye. Additionally, both the extent and the amount of stimulation or inhibition of neovascularization or regression of neovascularization can easily be monitored over time.

15 Finally, the rabbit is exposed to any test reagent, and therefore the health of the rabbit is an indication of toxicity of the test reagent.

25 The fourth assay measures angiogenesis in the chimeric mouse:human mouse model and is referred to as the chimeric mouse assay. The assay has been described in detail by others, and further has been described herein to measure angiogenesis, neovascularization, and regression of tumor tissues. See Yan, et al., J. Clin. Invest., 91:986-996 (1993). The chimeric mouse assay is a useful assay model for in vivo angiogenesis because the transplanted skin grafts closely resemble normal human skin histologically and neovascularization of whole tissue is occurring wherein actual human blood vessels are growing from the grafted human skin into the human tumor tissue on the surface of the grafted human skin. The

30

35

- 45 -

origin of the neovascularization into the human graft can be demonstrated by immunohistochemical staining of the neovasculature with human-specific endothelial cell markers.

5           As demonstrated herein, the chimeric mouse assay demonstrates regression of neovascularization based on both the amount and extent of regression of new vessel growth. Furthermore, it is easy to monitor effects on the growth of any tissue  
10           transplanted upon the grafted skin, such as a tumor tissue. Finally, the assay is useful because there is an internal control for toxicity in the assay system. The chimeric mouse is exposed to any test reagent, and therefore the health of the mouse is  
15           an indication of toxicity.

#### Examples

          The following examples relating to this invention are illustrative and should not, of  
20           course, be construed as specifically limiting the invention. Moreover, such variations of the invention, now known or later developed, which would be within the purview of one skilled in the art are to be considered to fall within the scope  
25           of the present invention hereinafter claimed.

#### 1. Preparation of Synthetic Peptides

          The linear and cyclic polypeptides listed in Table 1 were synthesized using standard solid-phase  
30           synthesis techniques as, for example, described by Merrifield, Adv. Enzymol., 32:221-96, (1969), and Fields, G.B. and Noble, R.L., Int. J. Peptide Protein Res., 35:161-214, (1990).

          Two grams (g) of BOC-Gly-D-Arg-Gly-Asp-Phe-  
35           Val-OMe (SEQ ID NO 1) were first dissolved in 60 milliliters (ml) of methanol to which was added 1.5 ml of 2 N sodium hydroxide solution to form an admixture. The admixture was then stirred for 3



- 46 -

hours at 20 degrees C (20C). After evaporation, the residue was taken up in water, acidified to pH 3 with diluted HCl and extracted with ethyl acetate. The extract was dried over Na<sub>2</sub>SO<sub>4</sub>, evaporated again and the resultant BOC-Gly-D-Arg-Gly-Asp-Phe-Val-OH (SEQ ID NO 2) was stirred at 20C for 2 hours with 20 ml of 2 N HCl in dioxane. The resultant admixture was evaporated to obtain H-Gly-D-Arg-Gly-Asp-Phe-Val-OH (SEQ ID NO 3) that was subsequently dissolved in a mixture of 1800 ml of dichloromethane and 200 ml of dimethylformamide (DMF) followed by cooling to 0C. Thereafter, 0.5 g of dicyclohexylcarbodiimide (DCCI), 0.3 g of 1-hydroxybenzotriazole (HOBt) and 0.23 ml of N-methylmorpholine were added successively with stirring.

The resultant admixture was stirred for a further 24 hours at 0C and then at 20C for another 48 hours. The solution was concentrated and treated with a mixed bed ion exchanger to free it from salts. After the resulting resin was removed by filtration, the clarified solution was evaporated and the residue was purified by chromatography resulting in the recovery of cyclo(-Gly-D-Arg-Gly-Asp-Phe-Val) (SEQ ID NO 4). The following peptides, listed in Table 1 using single letter code amino acid residue abbreviations and identified by a peptide number designation, were obtained analogously: cyclo(Arg-Gly-Asp-D-Phe-Val) (SEQ ID NO 5); cyclo(Arg-Ala-Asp-D-Phe-Val) (SEQ ID NO 6); cyclo(Arg-D-Ala-Asp-Phe-Val) (SEQ ID NO 9); cyclo(Arg-Gly-Asp-Phe-D-Val) (SEQ ID NO 7). A peptide designated as 66203, having an identical sequence to that of peptide 62184, only differed from the latter by containing the salt HCl rather than the TFA salt present in 62184. In inhibition of angiogenesis assays as described in Example 7 where the synthetic peptides were used, the 66203

- 47 -

peptide having HCl was slightly more effective in inhibiting angiogenesis than the identical peptide in TFA.

5

Table 1

	<u>Peptide No.</u>	<u>Amino Acid Sequence*</u>	<u>SEQ ID NO</u>
	62181	cyclo(GrGDFV)	4
	62184	cyclo(RGDfV)	5
10	62185	cyclo(RADfV)	6
	62187	cyclo(RGDFv)	7
	62880	YTAECKPQVTRGDVF	8
	62186	cyclo(RaDFV)	9
	62175	cyclo(ARGDfL)	10
15	62179	cyclo(GRGDfL)	11
	62411	TRQVVC DLGNPM	12
	62503	GVVRNNEALARLS	13
	62502	TDVNGDGRHDL	14

20 \* Lower case letters indicate a D-amino acid;  
capital letters indicate a L-amino acid.

25 A peptide designated as 69601, having an identical sequence to that of peptide 62185, only differed from the latter by containing the salt HCl rather than the TFA salt present in 62184.

30 The cyclic peptide c-RADfV (69601) has been shown to inhibit binding of fibrinogen to the integrin  $\alpha_v\beta_3$ , and not inhibit binding of fibrinogen to the integrins  $\alpha_{IIb}\beta_3$  or  $\alpha_5\beta_1$  (Pfaff, et al., J. Biol. Chem., 269:20233-20238, 1994). Thus the peptide c-RADfV is specific for  $\alpha_v\beta_3$ .

## 2. Monoclonal Antibodies

35 The monoclonal antibody LM609 secreted by hybridoma ATCC HB 9537 was produced using standard hybridoma methods by immunization with isolated  $\alpha_v\beta_3$  adsorbed onto Sepharose-lentil lectin beads. The  $\alpha_v\beta_3$  had been isolated from human melanoma cells

- 48 -

designated M21, and antibody was produced as described by Cheresh et al., J. Biol. Chem., 262:17703-17711 (1987). M21 cells were provided by Dr. D.L. Morton (University of California at Los Angeles, CA) and grown in suspension cultures in RPMI 1640 culture medium containing 2 mM L-glutamine, 50 mg/ml gentamicin sulfate and 10 % fetal calf serum.

Monoclonal antibody LM609 has been shown to specifically immunoreact with  $\alpha_v\beta_3$  complex, and not immunoreact with  $\alpha_v$  subunit, with  $\beta_3$  subunit, or with other integrins.

### 3. Characterization of the Tissue Distribution of $\alpha_v\beta_3$ Expression

#### A. Immunofluorescence with Anti-Integrin Receptor Antibodies

During wound healing, the basement membranes of blood vessels express several adhesive proteins, including von Willebrand factor, fibronectin, and fibrin. In addition, several members of the integrin family of adhesion receptors are expressed on the surface of cultured smooth muscle and endothelial cells. See, Cheresh, Proc. Natl. Acad. Sci., USA, 84:6471 (1987); Janat et al., J. Cell Physiol., 151:588 (1992); and Cheng et al., J. Cell Physiol., 139:275 (1989). Among these integrins is  $\alpha_v\beta_3$ , the endothelial cell receptor for von Willebrand factor, fibrinogen (fibrin), and fibronectin as described by Cheresh, Proc. Natl. Acad. Sci., USA, 84:6471 (1987). This integrin initiates a calcium-dependent signaling pathway leading to endothelial cell migration, and therefore appears to play a fundamental role in vascular cell biology as described by Leavelsey et al., J. Cell Biol., 121:163 (1993).

To investigate the expression of  $\alpha_v\beta_3$  during angiogenesis, human wound granulation tissue or

- 49 -

adjacent normal skin was obtained from consenting patients, washed with 1 ml of phosphate buffered saline and embedded in O.T.C medium (Tissue Tek). The embedded tissues were snap frozen in liquid nitrogen for approximately 30 to 45 seconds. Six micron thick sections were cut from the frozen blocks on a cryostat microtome for subsequent immunoperoxidase staining with antibodies specific for either  $\beta_3$  integrins ( $\alpha_v\beta_3$  or  $\alpha_{IIB}\beta_3$ ) or the  $\beta_1$  subfamily of integrins.

The results of the staining of normal human skin and wound granulation tissue are shown in Figures 1A-1D. Monoclonal antibodies AP3 and LM534, directed to  $\beta_3$  and  $\beta_1$  integrins, respectively, were used for immunohistochemical analysis of frozen sections. Experiments with tissue from four different human donors yielded identical results. The photomicrographs are shown at magnification of 300x.

The  $\alpha_v\beta_3$  integrin was abundantly expressed on blood vessels in granulation tissue (Figure 1B) but was not detectable in the dermis and epithelium of normal skin from the same donor (Figure 1A). In contrast,  $\beta_1$  integrins were abundantly expressed on blood vessels and stromal cells in both normal skin (Figure 1C) and granulation tissue (Figure 1D) and, as previously shown as described by Adams et al., Cell, 63:425 (1991), on the basal cells within the epithelium.

#### B. Immunofluorescence with Anti-Ligand Antibodies

Additional sections of the human normal skin and granulation tissues prepared above were also examined for the presence of the ligands for the  $\beta_3$  and  $\beta_1$  integrins, von Willebrand factor and laminin, respectively. Von Willebrand factor localized to the blood vessels in normal skin

- 50 -

(Figure 2A) and granulation tissue (Figure 2B), whereas laminin localized to all blood vessels as well as the epithelial basement membrane in both tissue preparations (Figures 2C and 2D).

5

C. Distribution Anti- $\alpha_v\beta_3$  Antibodies on Cancer Tissue

In addition to the above analyses, biopsies of cancer tissue from human patients were also examined for the presence and distribution of  $\alpha_v\beta_3$ . The tissues were prepared as described Example 1A with the exception that they were stained with monoclonal antibody LM609 prepared in Example 2 that is specific for the integrin receptor complex,  $\alpha_v\beta_3$ . In addition, tumors were also prepared for microscopic histological analysis by fixing representative examples of tumors in Bulins Fixative for 8 hours and serial sections cut and H&E stained.

The results of immunoperoxidase staining of bladder, colon breast and lung cancer tissues are shown in Figures 3A-3D, respectively.  $\alpha_v\beta_3$  is abundantly expressed only on the blood vessels present in the four cancer biopsies analyzed and not on any other cells present in the tissue.

The results described herein thus show that the  $\alpha_v\beta_3$  integrin receptor is selectively expressed in specific tissue types, namely granulated, metastatic tissues and other tissues in which angiogenesis is occurring and not normal tissue where the formation of new blood vessels has stopped. These tissues therefore provide an ideal target for therapeutic aspects of this invention.

35

- 51 -

4. Identification of  $\alpha_v\beta_3$ -Specific Synthetic  
Peptides Detected by a Ligand-Receptor Binding  
Assay

5 The synthetic peptides prepared in Example 1  
were screened by measuring their ability to  
antagonize  $\alpha_v\beta_3$  and  $\alpha_{IIB}\beta_3$  receptor binding activity  
in purified ligand-receptor binding assays. The  
method for these binding studies has been described  
by Barbas et al., Proc. Natl. Acad. Sci., USA,  
10 90:10003-10007 (1993), Smith et al., J. Biol.  
Chem., 265:11008-11013 (1990), and Pfaff et al., J.  
Biol. Chem., 269:20233-20238 (1994), the  
disclosures of which are hereby incorporated by  
reference.

15 Herein described is a method of identifying  
antagonists in a ligand-receptor binding assay in  
which the receptor is immobilized to a solid  
support and the ligand and antagonist are soluble.  
Also describes a ligand-receptor binding assay in  
20 which the ligand is immobilized to a solid support  
and the receptor and antagonists are soluble.

Briefly, selected purified integrins were  
separately immobilized in Titertek microtiter wells  
at a coating concentration of 50 nanograms (ng) per  
25 well. The purification of the receptors used in  
the ligand-receptor binding assays are well known  
in the art and are readily obtainable with methods  
familiar to one of ordinary skill in the art.  
After incubation for 18 hours at 4C, nonspecific  
30 binding sites on the plate were blocked with 10  
milligrams/milliliter (mg/ml) of bovine serum  
albumin (BSA) in Tris-buffered saline. For  
inhibition studies, various concentrations of  
selected peptides from Table 1 were tested for the  
35 ability to block the binding of  $^{125}\text{I}$ -vitronectin or  
 $^{125}\text{I}$ -fibrinogen to the integrin receptors,  $\alpha_v\beta_3$  and  
 $\alpha_{IIB}\beta_3$ . Although these ligands exhibit optimal  
binding for a particular integrin, vitronectin for

- 52 -

$\alpha_v\beta_3$  and fibrinogen for  $\alpha_{IIB}\beta_3$ , inhibition of binding studies using peptides to block the binding of fibrinogen to either receptor allowed for the accurate determination of the amount in micromoles (uM) of peptide necessary to half-maximally inhibit the binding of receptor to ligand. Radiolabeled ligands were used at concentrations of 1 nM and binding was challenged separately with unlabeled synthetic peptides.

Following a three hour incubation, free ligand was removed by washing and bound ligand was detected by gamma counting. The data from the assays where selected cyclic peptides listed in Table 1 were used to inhibit the binding of receptors and radiolabeled fibrinogen to separately immobilized  $\alpha_v\beta_3$  and  $\alpha_{IIB}\beta_3$  receptors were highly reproducible with the error between data points typically below 11%. The  $IC_{50}$  data in micromoles ( $IC_{50}$  uM) are expressed as the average of duplicate data points  $\pm$  the standard deviation as shown in Table 2.

Table 2

Peptide No.	$\alpha_v\beta_3$ ( $IC_{50}$ uM)	$\alpha_{IIB}\beta_3$ ( $IC_{50}$ uM)
62181	1.96 $\pm$ 0.62	14.95 $\pm$ 7.84
62184	0.05 $\pm$ 0.001	0.525 $\pm$ 0.10
62185	0.885 $\pm$ 0.16	100 $\pm$ 0.001
62187	0.05 $\pm$ 0.001	0.26 $\pm$ 0.056
62186	57.45 $\pm$ 7.84	100 $\pm$ 0.001
62175	1.05 $\pm$ 0.07	0.63 $\pm$ 0.18
62179	0.395 $\pm$ .21	0.055 $\pm$ 0.007

Thus, the RGD-containing or RGD-derivatized cyclized peptides 62181, 62184, 62185 and 62187, each having one D- amino acid residue, exhibited preferential inhibition of fibrinogen binding to the  $\alpha_v\beta_3$  receptor as measured by the lower

- 53 -

concentration of peptide required for half-maximal inhibition as compared to that for the  $\alpha_{IIB}\beta_3$  receptor. In contrast, the other RGD-containing or RGD-derivatized cyclic peptides, 62186, 62175 and 62179, were either not as effective in blocking fibrinogen binding to  $\alpha_v\beta_3$  or exhibited preferential inhibition of fibrinogen binding to  $\alpha_{IIB}\beta_3$  as compared to  $\alpha_v\beta_3$ . These results are consistent with those recently published by Pfaff, et al., J. Biol. Chem., 26920233-20238 (1994) in which the cyclic peptide RGDFV (wherein F indicates a D- amino acid residue) specifically inhibited binding of fibrinogen to the  $\alpha_v\beta_3$  integrin and not to the  $\alpha_{IIB}\beta_3$  or  $\alpha_5\beta_1$  integrins. Similar inhibition of binding assays were performed with linearized peptides having or lacking an RGD motif, the sequences of which were derived from the  $\alpha_v$  receptor subunit,  $\alpha_{IIB}$  receptor subunit or vitronectin ligand amino acid residue sequences. The sequences of the linear peptides, 62880 (VN-derived amino acid residues 35-49), 62411 ( $\alpha_v$ -derived amino acid residues 676-687); 62503 ( $\alpha_v$ -derived amino acid residues 655-667) and 62502 ( $\alpha_{IIB}$ -derived amino acid residues 296-306), are listed in Table 1. Each of these peptides were used in separate assays to inhibit the binding of either vitronectin (VN) or fibrinogen (FG) to either  $\alpha_{IIB}\beta_3$  or  $\alpha_v\beta_3$ . The  $IC_{50}$  data in micromoles ( $IC_{50}$  uM) of an individual assay for each experiment is shown in Table 3.

Table 3

Peptide No.	$\alpha_{IIB}\beta_3$ $IC_{50}$ (uM)		$\alpha_v\beta_3$ $IC_{50}$ (uM)	
	FG	VN	FG	VN
62880	4.2	0.98	<0.1	0.5
62411	>100	>100	>100	>100
62503	>100	>100	>100	>100
62502	90	5	>100	>100



The results of inhibition of ligand binding assays to selected integrin receptors with linearized peptides show that only peptide 62880 was effective at inhibiting the half-maximal binding of either FG or VN to  $\alpha_v\beta_3$ , as measured by the lower concentration of peptide required for half-maximal inhibition as compared to  $\alpha_{IIB}\beta_3$  receptor. None of the other linearized peptides were effective at blocking ligand binding to  $\alpha_v\beta_3$ , although peptide 62502 was effective at blocking VN binding to  $\alpha_{IIB}\beta_3$ .

Thus, the ligand-receptor assay described herein can be used to screen for both circular or linearized synthetic peptides that exhibit selective specificity for a particular integrin receptor, specifically  $\alpha_v\beta_3$ , as used as vitronectin receptor ( $\alpha_v\beta_3$ ) antagonists in practicing this invention.

## 5. Characterization of the Untreated Chick Chorioallantoic Membrane (CAM)

### A. Preparation of the CAM

Angiogenesis can be induced on the chick chorioallantoic membrane (CAM) after normal embryonic angiogenesis has resulted in the formation of mature blood vessels. Angiogenesis has been shown to be induced in response to specific cytokines or tumor fragments as described by Leibovich et al., Nature, 329:630 (1987) and Ausprunk et al., Am. J. Pathol., 79:597 (1975). CAMs were prepared from chick embryos for subsequent induction of angiogenesis and inhibition thereof as described in Examples 6 and 7, respectively. Ten day old chick embryos were obtained from McIntyre Poultry (Lakeside, CA) and incubated at 37C with 60% humidity. A small hole was made through the shell at the end of the egg

- 55 -

directly over the air sac with the use of a small  
crafts drill (Dremel, Division of Emerson Electric  
Co. Racine WI). A second hole was drilled on the  
broad side of the egg in a region devoid of  
embryonic blood vessels determined previously by  
candling the egg. Negative pressure was applied to  
the original hole, which resulted in the CAM  
(chorioallantoic membrane) pulling away from the  
shell membrane and creating a false air sac over  
the CAM. A 1.0 centimeter (cm) x 1.0 cm square  
window was cut through the shell over the dropped  
CAM with the use of a small model grinding wheel  
(Dremel). The small window allowed direct access  
to the underlying CAM.

The resultant CAM preparation was then either  
used at 6 days of embryogenesis, a stage marked by  
active neovascularization, without additional  
treatment to the CAM reflecting the model used for  
evaluating effects on embryonic neovascularization  
or used at 10 days of embryogenesis where  
angiogenesis has subsided. The latter preparation  
was thus used in this invention for inducing  
renewed angiogenesis in response to cytokine  
treatment or tumor contact as described in Example  
6.

#### B. Histology of the CAM

To analyze the microscopic structure of  
the chick embryo CAMs and/or human tumors that were  
resected from the chick embryos as described in  
Example 8, the CAMs and tumors were prepared for  
frozen sectioning as described in Example 3A. Six  
micron thick sections were cut from the frozen  
blocks on a cryostat microtome for  
immunofluorescence analysis.

Figure 4 shows a typical photomicrograph of an  
area devoid of blood vessels in an untreated 10 day  
old CAM. As angiogenesis in the CAM system is

- 56 -

subsiding by this stage of embryogenesis, the system is useful in this invention for stimulating the production of new vasculature from existing vessels from adjacent areas into areas of the CAM currently lacking any vessels.

C. Integrin Profiles in the CAM Detected by Immunofluorescence

To view the tissue distribution of integrin receptors present in CAM tissues, 6 micron (um) frozen sections of both tumor tissue and chick embryo CAM tissues were fixed in acetone for 30 seconds and stained by immunofluorescence with 10 micrograms/milliliter (ug/ml) mAb CSAT, a monoclonal antibody specific for the  $\beta_1$  integrin subunit as described by Buck et al., J. Cell Biol., 107:2351 (1988) and thus used for controls, or LM609 as prepared in Example 2. Primary staining was followed by staining with a 1:250 dilution of goat anti-mouse rhodamine labeled secondary antibody (Tago) to allow for the detection of the primary immunoreaction product. The sections were then analyzed with a Zeiss immunofluorescence compound microscope.

The results of the immunofluorescence analysis show that the mature blood vessels present in an untreated 10 day chick embryo expressed the integrin  $\beta_1$  subunit (Figure 5A). In contrast, in a serial section of the tissue shown in Figure 5A, no immunoreactivity with LM609 was revealed (Figure 5B). Thus, the integrin  $\alpha_v\beta_3$  detected by the LM609 antibody was not actively being expressed by the mature blood vessels present in a 10 day old untreated chick embryo. As shown in the CAM model and in the following Examples, while the blood vessels are undergoing new growth in normal embryogenesis or induced by either cytokines or tumors, the blood vessels are expressing  $\alpha_v\beta_3$ .

- 57 -

However, following active neovascularization, once the vessels have stopped developing, the expression of  $\alpha_v\beta_3$  diminishes to levels not detectable by immunofluorescence analysis. This regulation of  $\alpha_v\beta_3$  expression in blood vessels undergoing angiogenesis as contrasted to the lack of expression in mature vessels provides for the unique ability of this invention to control and inhibit angiogenesis as shown in the following Examples as modeled using the CAM angiogenesis assay system.

#### 6. CAM Angiogenesis Assay

##### 15 A. Angiogenesis Induced by Growth Factors

Angiogenesis has been shown to be induced by cytokines or growth factors as referenced in Example 5A. In the experiments described herein, angiogenesis in the CAM preparation described in Example 5 was induced by growth factors that were topically onto the CAM blood vessels as described herein.

Angiogenesis was induced by placing a 5 millimeter (mm) X 5 mm Whatman filter disk (Whatman Filter paper No.1) saturated with Hanks Balanced Salt Solution (HBSS, GIBCO, Grand Island, NY) or HBSS containing 150 nanograms/milliliter (ng/ml) recombinant basic fibroblast growth factor ( $\beta$ FGF) (Genzyme, Cambridge, MA) on the CAM of a 10-day chick embryo in a region devoid of blood vessels and the windows were latter sealed with tape. In other assays, 125 ng/ml  $\beta$ FGF was also effective at inducing blood vessel growth. Angiogenesis was monitored by photomicroscopy after 72 hours. CAMs were snap frozen, and 6  $\mu$ m cryostat sections were fixed with acetone and stained by immunofluorescence as described in Example 5C with

- 58 -

10 ug/ml of either anti- $\beta_1$  monoclonal antibody CSAT or LM609.

5 The immunofluorescence photomicrograph in Figure 5C shows enhanced expression of  $\alpha_v\beta_3$  during  $\beta$ FGF-induced angiogenesis on the chick CAM in contrast with the absence of  $\alpha_v\beta_3$  expression in an untreated chick CAM as shown in Figure 5B.  $\alpha_v\beta_3$  was readily detectable on many (75% to 80%) of the vessels on the  $\beta$ FGF-treated CAMs. In addition, the expression of integrin  $\beta_1$  did not change from that seen in an untreated CAM as  $\beta_1$  was also readily detectable on stimulated blood vessels.

10 The relative expression of  $\alpha_v\beta_3$  and  $\beta_1$  integrins was then quantified during  $\beta$ FGF-induced angiogenesis by laser confocal image analysis of the CAM cryostat sections. The stained sections were then analyzed with a Zeiss laser confocal microscope. Twenty-five vessels stained with LM609 and 15 stained with CSAT (average size ~ 1200 sq mm<sup>2</sup>, range 350 to 3,500 mm<sup>2</sup>) were selected from random fields and the average rhodamine fluorescence for each vessel per unit area was measured in arbitrary units by laser confocal image analysis. Data are expressed as the mean fluorescence intensity in arbitrary units of vessels  $\pm$  standard error (SE).

15 The results plotted in Figure 6 show that staining of  $\alpha_v\beta_3$  was significantly enhanced (four times higher) on CAMs treated with  $\beta$ FGF as determined by the Wilcoxon Rank Sum Test ( $P < 0.0001$ ) whereas  $\beta_1$  staining was not significantly different with  $\beta$ FGF treatment.

20 The CAM assay was further used to examine the effect of another potent angiogenesis inducer, tumor necrosis factor-alpha (TNF $\alpha$ ), on the expression of  $\beta_1$  and  $\beta_3$  integrins. Filter disks impregnated with either  $\beta$ FGF or TNF $\alpha$  and placed on

- 59 -

CAMs from 10 day embryos were found to promote local angiogenesis after 72 hours.

The results are shown in the photomicrographs of CAMs either untreated (Figure 7A), treated with  $\beta$ FGF (Figure 7B) or treated with  $\text{TNF}\alpha$  (Figure 7C). Blood vessels are readily apparent in both the  $\beta$ FGF and  $\text{TNF}\alpha$  treated preparations but are not present in the untreated CAM. Thus, the topical application of a growth factor/cytokine resulted in the induction of angiogenesis from mature vessels in an adjacent area into the area originally devoid of blood vessels. In view of the  $\beta$ FGF-induced blood vessels and concomitant expression of  $\alpha_v\beta_3$ , as shown in Figure 5C, treatment of  $\text{TNF}\alpha$  results in comparable activities.

These findings indicate that in both human and chick, blood vessels involved in angiogenesis show enhanced expression of  $\alpha_v\beta_3$ . Consistent with this, expression of  $\alpha_v\beta_3$  on cultured endothelial cells can be induced by various cytokines in vitro as described by Janat et al., J. Cell Physiol., 151:588 (1992); Enenstein et al., Exp. Cell Res., 203:499 (1992) and Swerlick et al., J. Invest. Derm., 99:715 (1993).

The effect on growth-factor induced angiogenesis by antibody and peptide inhibitors is presented in Examples 7A and 7B.

#### B. Embryonic Angiogenesis

The CAM preparation for evaluating the effect of angiogenesis inhibitors on the natural formation of embryonic neovasculature was the 6 day embryonic chick embryo as previously described. At this stage in development, the blood vessels are undergoing de novo growth and thus provides a useful system for determining if  $\alpha_v\beta_3$  participates in embryonic angiogenesis. The CAM system was prepared as described above with the exception that

- 60 -

the assay was performed at embryonic day 6 rather than at day 10. The effect on embryonic angiogenesis by treatment with antibodies and peptides of this invention are presented in Example 7C.

C. Angiogenesis Induced by Tumors

To investigate the role of  $\alpha_v\beta_3$  in tumor-induced angiogenesis, various  $\alpha_v\beta_3$ -negative human melanoma and carcinoma fragments were used in the CAM assay that were previously grown and isolated from the CAM of 17-day chick embryo as described by Brooks et al., J. Cell Biol., 122:1351 (1993) and as described herein. The fragments induced extensive neovascularization in the presence of buffer alone.

Angiogenesis was induced in the CAM assay system by direct apposition of a tumor fragment on the CAM. Preparation of the chick embryo CAM was identical to the procedure described above. Instead of a filter paper disk, a 50 milligram (mg) to 55 mg in weight fragment of one of human melanoma tumor M21L, human lung carcinoma tumor UCLAP-3, human pancreatic carcinoma cell line FG (Cheresh et al., Cell 58:945-953, 1989), or human laryngeal carcinoma cell line HEp3, all of which are  $\alpha_v\beta_3$  negative tumors, was placed on the CAM in an area originally devoid of blood vessels.

The M21L human melanoma cell line, UCLAP-3 human lung carcinoma cell line, FG pancreatic carcinoma cell line, or HEp3 human laryngeal carcinoma cell line, all  $\alpha_v\beta_3$  negative, were used to grow the solid human tumors on the CAMs of chick embryos. A single cell suspension of  $8 \times 10^6$  M21-L, UCLAP-3, and FB or  $5 \times 10^5$  HEp3 cells were first applied to the CAMs in a total volume of 30 microliters (ul) of sterile HBSS. The windows were sealed with tape and the embryos were incubated for

- 61 -

7 days to allow growth of human tumor lesions. At the end of 7 days, now a 17-day embryo, the tumors were resected from the CAMs and trimmed free of surrounding CAM tissue. The tumors were sliced  
5 into 50 mg to 55 mg tumor fragments for use in either angiogenesis or tumor growth assays. The tumor fragments were placed on a new set of 10 day chick embryo CAMs as described in Example 6A in an area devoid of blood vessels.

10 Tumors grown in vivo on the chick embryo CAMs were stained for  $\alpha_v\beta_3$  expression with mAb LM609 as described in Example 3A. No specific staining of tumor cells was observed indicating a lack of  $\alpha_v\beta_3$  expression.

15 These CAM tumor preparations were then subsequently treated as described in Examples 7D and 7E for measuring the effects of antibodies and peptides on tumor-induced angiogenesis. The CAM tumor preparations were also treated as described  
20 in Examples 8, 9, and 12 for measuring the effects of antibodies and peptides on regression of tumors and apoptosis of angiogenic blood vessels and vascular cells.

25 7. Inhibition of Angiogenesis as Measured in the CAM Assay

A. Inhibition of Growth Factor-Induced Angiogenesis by Topical Application of Inhibitors

30 1) Treatment with Monoclonal Antibodies

To determine whether  $\alpha_v\beta_3$  plays an active role in angiogenesis, filter disks saturated with  $\beta$ FGF or TNF $\alpha$  were placed on CAMs then the  
35 monoclonal antibodies (also referred to as mAb), LM609 (specific for  $\alpha_v\beta_3$ ), CSAT (specific for  $\beta_1$ ), or P3G2 (specific for  $\alpha_v\beta_5$ ) were added to the preparation.



- 62 -

Angiogenesis was induced on CAMs from 10 day chick embryos by filter disks saturated with  $\beta$ FGF. Disks were then treated with 50  $\mu$ l HBSS containing 25  $\mu$ g of mAb in a total volume of 25  $\mu$ l of sterile HBSS at 0, 24, and 48 hours. At 72 hours, CAMs were harvested and placed in a 35 mm petri dish and washed once with 1 ml of phosphate buffered saline. The bottom side of the filter paper and CAM tissue was then analyzed under an Olympus stereo microscope, with two observers in a double-blind fashion. Angiogenesis inhibition was considered significant when CAMs exhibited >50% reduction in blood vessel infiltration of the CAM directly under the disk. Experiments were repeated four times per antibody, with 6 to 7 embryos per condition.

The results of the effects of mAb treatment on  $\beta$ FGF-induced angiogenesis is shown in Figures 8A-8E. An untreated CAM preparation devoid of blood vessels is shown in Figure 8A to provide a comparison with the  $\beta$ FGF-blood vessel induction shown in Figure 8B and effects thereon by the mAbs in Figures 8C-8E. About 75% of these CAMs treated with mAb LM609 exhibited >50% inhibition of angiogenesis as shown in Figure 8E, and many of these appeared devoid of vessel infiltration. In contrast, the buffer control (Figure 8A) and disks treated with mAbs CSAT (Figure 8C) and P3G2 (Figure 8D) consistently showed extensive vascularization.

Identical results were obtained when angiogenesis was induced with  $\text{TNF}\alpha$ . To examine the effects of these same antibodies on preexisting mature blood vessels present from normal vessel development adjacent to the areas devoid of vessels, filter disks saturated with mAbs were placed on vascularized regions of CAMs from 10 day embryos that did not receive topical application of cytokine. None of the three mAbs affected preexisting vessels, as assessed by visualization

- 63 -

under a stereo microscope. Thus, mAb LM609 selectively inhibited only new blood vessel growth and did not effect mature blood vessels present in adjacent areas. This same effect was seen with the application of synthetic peptides either applied topically or intravenously as described in Examples 7A2) and 7E2), respectively.

## 2) Treatment with Synthetic Peptides

CAM assays were also performed with the synthetic peptides of this invention to determine the effect of cyclic and linearized peptides on growth factor induced angiogenesis. The peptides were prepared as described in Example 1 and 80 ug of peptide was presented in a total volume of 25 ul of sterile HBSS. The peptide solution was applied to the CAM preparation immediately and then again at 24 and 48 hrs. At 72 hours the filter paper and surrounding CAM tissue was dissected and viewed as described above.

Results from this assay revealed were similar to those shown in Figures 9A-9C as described in Example 7E2) where synthetic peptides were intravenously injected into tumor induced blood vessels. Here, with the control peptide, 62186, the  $\beta$ FGF-induced blood vessels remained undisturbed as shown in Figure 9A. In contrast when the cyclic RGD peptide, 62814, was applied to the filter, the formation of blood vessels was inhibited leaving the area devoid of new vasculature. This effect was similar in appearance to that shown in Figure 9B as described in Example 7E2) below. In addition, also as shown in Figure 9C for intravenously injected peptides, in areas in which mature blood vessels were present yet distant from the placement of the growth-factor saturated filter, no effect was seen with the topical treatment of synthetic peptides on these outlying

- 64 -

vessels. The inhibitory activity of the peptides on angiogenesis thus is limited to the areas of angiogenesis induced by growth factors and does not effect adjacent preexisting mature vessels or result in any deleterious cytotoxicity to the surrounding area.

Similar assays are performed with the other synthetic peptides prepared in Example 1 and listed in Table 1.

B. Inhibition of Growth Factor-Induced Angiogenesis by Intravenous Application of Inhibitors

1) Treatment with Monoclonal Antibodies

The effect on growth factor-induced angiogenesis with monoclonal antibodies intravenously injected into the CAM preparation was also evaluated for use in this invention.

The preparation of the chick embryo CAMs for intravenous injections were essentially as described in Example 7A with some modifications. During the candling procedures prominent blood vessels were selected and marks were made on the egg shell to indicate their positions. The holes were drilled in the shell and the CAMs were dropped and  $\beta$ FGF saturated filter papers were placed on the CAMs as described above. The windows were sealed with sterile tape and the embryos were replaced in the incubator. Twenty four hours later, a second small window was carefully cut on the lateral side of the egg shell directly over prominent blood vessels selected previously. The outer egg shell was carefully removed leaving the embryonic membranes intact. The shell membrane was made transparent with a small drop of mineral oil

(Perkin-Elmer Corp, Norwalk, CT) which allowed the blood vessels to be visualized easily. Purified sterile mAbs, or synthetic peptides, the latter of

- 65 -

which are described below, were inoculated directly into the blood vessels once with a 30 gauge needle at a dose of 200 ug of IgG per embryo in a total volume of 100 ul of sterile PBS. The windows were sealed with tape and the embryos were allowed to incubate until 72 hours. The filter disks and surrounding CAM tissues were analyzed as described before.

To determine the localization of LM609 mAb in CAM tissues or in tumor tissues, as shown herein and in the following Examples, that were previously inoculated intravenously with LM609, the fixed sections were blocked with 2.5% BSA in HBSS for 1 hour at room temperature followed by staining with a 1:250 dilution of goat anti-mouse rhodamine labeled secondary antibody (Tago). The sections were then analyzed with a Zeiss immunofluorescence compound microscope.

The results of intravenous antibody treatment to the  $\beta$ FGF induced blood vessel CAM preparation are shown in Figures 10A-10C. In Figure 10A, angiogenesis induced as a result of  $\beta$ FGF treatment is shown. No change to the presence of  $\beta$ FGF induced vasculature was seen with intravenous exposure to mAb P3G2, an anti- $\alpha_v\beta_3$  antibody, as shown in Figure 10B. In contrast, treatment of the  $\beta$ FGF induced angiogenesis CAM preparation with LM609, an anti- $\alpha_v\beta_3$  antibody, resulted in the complete inhibition of growth of new vessels into the filter area as shown in Figure 10C. The inhibitory effect on angiogenesis is thus resulting from the inhibition of  $\alpha_v\beta_3$  receptor activity by the LM609 anti- $\alpha_v\beta_3$ -specific antibody. Since the blocking of the  $\alpha_v\beta_3$  does not inhibit the formation of neovasculature into the CAMs filter site,  $\alpha_v\beta_3$  thus is not essential as compared to  $\alpha_v\beta_3$  for growth of new vessels.

- 66 -

2) Treatment with Synthetic Peptides

The synthetic peptides prepared in Example 1 are separately intravenously injected into the growth factor induced blood vessels in the CAM preparation as described above. The effect of the peptides on the viability of the vessels is similarly assessed.

C. Inhibition of Embryonic Angiogenesis by Topical Application1) Treatment with Monoclonal Antibodies

To determine whether  $\alpha_v\beta_3$  participates in embryonic angiogenesis, the effect of LM609 on de novo growth of blood vessels on CAMs was examined in 6 day embryos, a stage marked by active neovascularization as described in Example 5A. The CAM assay was prepared as described in Example 6C with the subsequent topical application of disks saturated with mAbs placed on CAMs of 6 day old embryos in the absence of cytokines. After 3 days, CAMs were resected and photographed. Each experiment included 6 embryos per group and was repeated 2 times.

Antibody LM609 (Figure 11C), but not CSAT (Figure 11A) or P3G2 (Figure 11B), prevented vascular growth under these conditions; this indicates that  $\alpha_v\beta_3$  plays a substantial role in embryonic neovascularization that was independent of added growth factors for induction of angiogenesis.

2) Treatment with Synthetic Peptides

The synthetic peptides prepared in Example 1 are separately added to the embryonic CAM preparation prepared above and as described in Example 5A2) by either topical application to the CAM or intravenous application to blood vessels.

- 67 -

The effect of the peptides on the viability of the vessels is similarly assessed.

D. Inhibition of Tumor-Induced Angiogenesis  
by Topical Application

1) Treatment with Monoclonal Antibodies

In addition to the angiogenesis assays described above where the effects of anti- $\alpha_v\beta_3$  antagonists, LM609 and peptides 62181, 62184, 62185, 62187 and 62880, on embryonic angiogenesis were evaluated, the role of  $\alpha_v\beta_3$  in tumor-induced angiogenesis was also investigated. As an inducer,  $\alpha_v\beta_3$ -negative human M21-L melanoma fragments previously grown and isolated from the CAM of a 17-day chick embryo were used. The fragments were prepared as described in Example 6C.

As described above in Example 7A1), mAbs were separately topically applied to the tumor fragments at a concentration of 25 ug in 25 ul of HBSS and the windows were then sealed with tape. The mAbs were added again in the same fashion at 24 hours and 48 hours. At 72 hours, the tumors and surrounding CAM tissues were analyzed as described above in Example 7A1).

As described in Example 6C, tumors were initially derived by transplanting cultured M21-L cells, which do not to express integrin  $\alpha_v\beta_3$ , as described by Felding-Habermann et al., J. Clin. Invest., 89:2018 (1992) onto the CAMs of 10-day old chick embryos. These  $\alpha_v\beta_3$ -negative fragments induced extensive neovascularization in the presence of buffer alone, or mAbs CSAT (anti- $\beta_1$ ) or P3G2 (anti- $\alpha_v\beta_5$ ). In contrast, mAb LM609 (anti- $\alpha_v\beta_3$ ) abolished the infiltration of most vessels into the tumor mass and surrounding CAM.

In order to quantitate the effect of the mAbs on the tumor-induced angiogenesis, blood vessels entering the tumor within the focal plane of the

- 68 -

CAM were counted under a stereo microscope by two observers in a double-blind fashion. Each data bar presented in Figure 12 represents the mean number of vessels  $\pm$  SE from 12 CAMs in each group representing duplicate experiments.

This quantitative analysis revealed a three-fold reduction in the number of vessels entering tumors treated with mAb LM609 compared to tumors treated with buffer or the other mAbs, P3G2 or CSAT ( $P < 0.0001$ ) as determined by Wilcoxon Rank Sum Test. The fact that M21-L tumors do not express  $\alpha_v\beta_3$  indicates that mAb LM609 inhibits angiogenesis by directly affecting blood vessels rather than the tumor cells. These results correspond with the histological distribution of  $\alpha_v\beta_3$  in cancer tissue biopsies shown in Figure 3A-3D where the distribution of  $\alpha_v\beta_3$  was limited to the blood vessels in the tumor and not to the tumor cells themselves.

## 2) Treatment with Synthetic Peptides

The synthetic peptides prepared in Example 1 are topically applied to the tumor-induced angiogenic CAM assay system as described above. The effect of the peptides on the viability of the vessels is similarly assessed.

## E. Inhibition of Tumor-Induced Angiogenesis by Intravenous Application

### 1) Treatment with Monoclonal Antibodies

Tumor-induced blood vessels prepared as described in Example 7D1) were also treated with mAbs applied by intravenous injection. Tumors were placed on the CAMs as described in Example 7D1) and the windows sealed with tape and 24 hours latter, 200 ug of purified mAbs were inoculated once intravenously in chick embryo blood vessels as described previously. The chick embryos were then

- 69 -

allowed to incubate for 7 days. The extent of angiogenesis was then observed as described in above. As described in Example 8 below, after this time period, the tumors were resected and analyzed by their weight to determine the effect of antibody exposure on tumor growth or suppression.

## 2) Treatment with Synthetic Peptides

The effects of peptide exposure to tumor-induced vasculature in the CAM assay system was also assessed. The tumor-CAM preparation was used as described above with the exception that instead of intravenous injection of a mAb, synthetic peptides prepared as described in Example 1 and Example 7A2) were separately intravenously injected into visible blood vessels.

The results of CAM assays with the cyclic peptide, 66203 containing the HCl salt, and control peptide, 62186, are shown in Figures 9A-9C. In Figure 9A, the treatment with the control peptide did not effect the abundant large blood vessels that were induced by the tumor treatment to grow into an area originally devoid of blood vessels of the CAM. In contrast when the cyclic RGD peptide, 66203, an antagonist to  $\alpha_v\beta_3$ , was applied to the filter, the formation of blood vessels was inhibited leaving the area devoid of new vasculature as shown in Figure 9B. The inhibitory effect of the RGD-containing peptide was specific and localized as evidenced by an absence of any deleterious effects to vessels located adjacent to the tumor placement. Thus, in Figure 9C, when inhibitory peptides are intravenously injected into the CAM assay system, no effect was seen on the preexisting mature vessels present in the CAM in areas adjacent yet distant from the placement of the tumor. The preexisting vessels in this location were not affected by the inhibitory



- 70 -

peptide that flowed within those vessels although the generation of new vessels from these preexisting vessels into the tumor mass was inhibited. Thus, synthetic peptides including 66203 and 62184, previously shown in ligand-receptor assays in Example 4 to be antagonists of  $\alpha_v\beta_3$ , have now been demonstrated to inhibit angiogenesis that is limited to vessels undergoing development and not to mature preexisting vessels. In addition, the intravenous infusion of peptides does not result in any deleterious cytotoxicity to the surrounding area as evidenced by the intact vasculature in Figure 9C.

Similar assays are performed with the other synthetic peptides prepared in Example 1 and listed in Table 1.

8. Inhibition of Tumor Tissue Growth With  $\alpha_v\beta_3$  Antagonists As Measured in the CAM Assay

As described in Example 7D1), in addition to visually assessing the effect of anti- $\alpha_v\beta_3$  antagonists on growth factor or tumor induced angiogenesis, the effect of the antagonists was also assessed by measuring any changes to the tumor mass following exposure. For this analysis, the tumor-induced angiogenesis CAM assay system was prepared as described in Example 6C and 7D. At the end of the 7 day incubation period, the resulting tumors were resected from the CAMs and trimmed free of any residual CAM tissue, washed with 1 ml of phosphate buffer saline and wet weights were determined for each tumor.

In addition, preparation of the tumor for microscopic histological analysis included fixing representative examples of tumors in Bouin's Fixative for 8 hours and embedding in paraffin. Serial sections were cut and stained with hematoxylin and eosin (H&E) for microscopic

analysis. Gladson, et al., J. Clin. Invest., 88:1924 (1991). Sections were photographed with an Olympus compound microscope at 250x.

5                    A.    Topical Application

The results of typical human melanoma tumor (M21L) weights resulting from topical application of control buffer (HBSS), P3G2 (anti- $\alpha_v\beta_5$ ) or LM609 (anti- $\alpha_v\beta_3$ ) are listed in Table 4. A number of embryos were evaluated for each treatment with the average tumor weight in milligrams (mg) from each being calculated along with the SE of the mean as shown at the bottom of the table.

15 Table 4

	<u>Embryo No.</u>	<u>mAb Treatment</u>	<u>Tumor Weight</u>
	<u>(mg)</u>		
20	1	HBSS	108
	2		152
	3		216
	4		270
	5		109
	6		174
25	1	P3G2	134
	2		144
	3		408
	4		157
	5		198
	6		102
	7		124
	8		99
30	1		
	2		
	3		
	4		
35	1	LM609	24
	2		135
	3		17
	4		27

- 72 -

5	35
6	68
7	48
8	59

5

<u>mAb Treatment</u>	<u>Average Tumor Weight (mg)</u>
HBSS control	172 $\pm$ 26
P3G2	171 $\pm$ 36
LM609	52 $\pm$ 13

10

Exposure of a  $\alpha_v\beta_3$ -negative human melanoma tumor mass in the CAM assay system to LM609 caused the decrease of the untreated average tumor weight of 172 mg  $\pm$  26 to 52 mg  $\pm$  13. The P3G2 antibody had no effect on the tumor mass. Thus, the blocking of the  $\alpha_v\beta_3$  receptor by the topical application of  $\alpha_v\beta_3$ -specific LM609 antibody resulted in a regression of tumor mass along with an inhibition of angiogenesis as shown in the preceding Examples. The measured diameter of the tumor mass resulting from exposure to P3G2 was approximately 8 millimeters to 1 centimeter on average. In contrast, the LM609-treated tumors were on average 2 to 3 millimeters in diameter.

15

20

25

Frozen sections of these tumors revealed an intact tumor cytoarchitecture for the tumor exposed to P3G2 in contrast to a lack of organized cellular structure in the tumor exposed to LM609.  $\alpha_v\beta_3$  receptor activity is therefore essential for an  $\alpha_v\beta_3$  negative tumor to maintain its mass nourished by development of  $\alpha_v\beta_3$ -expressing neovasculature. The blocking of  $\alpha_v\beta_3$  with the  $\alpha_v\beta_3$  antagonists of this invention results in the inhibition of angiogenesis into the tumor ultimately resulting in the diminution of tumor mass.

30

35

#### B. Intravenous Application

- 73 -

The results of typical carcinoma tumor (UCLAP-3) weights resulting from intravenous application of control buffer (PBS, phosphate buffered saline), CSAT (anti- $\beta_1$ ) or LM609 (anti- $\alpha_v\beta_3$ ) are listed in Table 5. A number of embryos were evaluated for each treatment with the average tumor weight from each being calculated along with the SE of the mean as shown at the bottom of the table.

Table 5

	<u>Embryo No.</u> <u>(mg)</u>	<u>mAb Treatment</u>	<u>Tumor Weight</u>
15	1	PBS	101
	2		80
	3		67
	4		90
20	1	CSAT	151
	2		92
	3		168
	4		61
	5		70
25	1	LM609	16
	2		54
	3		30
	4		20
30	5		37
	6		39
	7		12
35		<u>mAb Treatment</u>	<u>Average Tumor Weight (mg)</u>
		PBS control	85 $\pm$ 7
		CSAT	108 $\pm$ 22
		LM609	30 $\pm$ 6

- 74 -

Exposure of a  $\alpha_v\beta_3$ -negative human carcinoma tumor mass in the CAM assay system to LM609 caused the decrease of the untreated average tumor weight of  $85 \text{ mg} \pm 7$  to  $30 \text{ mg} \pm 6$ . The CSAT antibody did not significantly effect the weight of the tumor mass. Thus, the blocking of the  $\alpha_v\beta_3$  receptor by the intravenous application of  $\alpha_v\beta_3$ -specific LM609 antibody resulted in a regression of a carcinoma as it did for the melanoma tumor mass above along with an inhibition of angiogenesis as shown in the preceding Examples. In addition, human melanoma tumor growth was similarly inhibited by intravenous injection of LM609.

9. Regression of Tumor Tissue Growth With  $\alpha_v\beta_3$  Antagonists As Measured in the CAM Assay

To assess the effects of  $\alpha_v\beta_3$  antagonists on tumor growth and survival, fragments of human melanoma and fragments of carcinomas of the lung, pancreas, and larynx were placed on CAMS of 10-day old embryos as described in Example 5A.

A. Intravenous Application

- 1) Treatment with Monoclonal Antibodies
  - a. Treatment with LM609 (anti- $\alpha_v\beta_3$ ) and CSAT (anti- $\beta_1$ )

Twenty four hours after implantation of CAM with carcinoma fragments of  $\alpha_v\beta_3$ -negative human melanoma M21-L, pancreatic carcinoma FG, human lung carcinoma UCLAP-3, or human laryngeal carcinoma HEp3, embryos were injected intravenously with PBS alone or a single dose ( $300 \text{ ug}/100 \text{ ul}$ ) of either mAb LM609 (anti- $\alpha_v\beta_3$ ) or CSAT (anti- $\beta_1$ ). Tumors were allowed to propagate for six additional days. At the end of the incubation period the tumors were carefully resected and trimmed free of surrounding CAM

- 75 -

tissue. Tumor resections were performed by two independent investigators removing only the easily definable solid tumor mass. The tumors had well defined margins, thus the thin semi-transparent membrane (CAM) which is readily distinguishable from the solid tumor mass was removed without disturbing the tumor mass itself. The resected tumors were weighed and examined morphologically and histologically.

As shown in Figure 13, wet tumor weights at the end of 7 days were determined and compared to initial tumor weights before treatments. Each bar represents the mean  $\pm$  S.E. of 5-10 tumors per group. mAb LM609 inhibited tumor growth significantly ( $p < 0.001$ ) as compared to controls in all tumors tested. Tumors treated with PBS or CSAT proliferated in all cases. In contrast, mAb LM609 not only prevented the growth of these tumors but induced extensive regression in most cases. Importantly, these tumor cells do not express integrin  $\alpha_v\beta_3$ , demonstrating that the inhibition of growth was due to the anti-angiogenic effects of this antibody on neovasculature rather than upon the tumor cells directly.

b. Treatment with LM609 (anti- $\alpha_v\beta_3$ ) and P3G2 (anti- $\alpha_v\beta_5$ )

Human M21-L melanoma tumor fragments (50 mg) were implanted on the CAMs of 10 day old embryos as described in Example 5A. Twenty four hours later, embryos were injected intravenously with PBS alone or a single dose (300 ug/100 ul) of either mAb LM609 (anti- $\alpha_v\beta_3$ ) or P3G2 (anti- $\alpha_v\beta_5$ ). Tumors were allowed to propagate as described in Example 9A1) above and were examined morphologically and histologically as herein described.

- 76 -

Representative examples of M21-L tumors treated with mAbs P3G2 (anti- $\alpha_v\beta_3$ ) or LM609 (anti- $\alpha_v\beta_3$ ) were examined morphologically. The P3G2-treated tumors were large (8 mm in diameter) and well vascularized whereas those treated with mAb LM609 were much smaller (3 mm in diameter) and lacked detectable blood vessels.

The tumors were further examined by the preparation of histological sections and staining with hematoxylin and eosin as described in Example 9A1)a. As shown in Figure 14 (upper panel), tumors treated with mAb P3G2 (anti- $\alpha_v\beta_3$ ) showed numerous viable and actively dividing tumor cells as indicated by mitotic figures (arrowheads) as well as by multiple blood vessels (arrows) throughout the tumor stroma. In contrast, few if any viable tumor cells or blood vessels were detected in tumors treated with mAb LM609 (anti- $\alpha_v\beta_3$ ) (Figure 14, lower panel). These results demonstrate that antagonists of integrin  $\alpha_v\beta_3$  inhibit tumor-induced angiogenesis leading to the growth arrest and regression of a variety of human tumors *in vivo*. It is important to point out that embryos examined after seven days of tumor growth (embryonic day 17) appeared normal upon gross examination whether or not they were treated with an  $\alpha_v\beta_3$  antagonist. These findings indicate that antagonists of this integrin appear non-toxic to the developing embryos.

## 2) Treatment With Synthetic Peptides

Human M21-L melanoma tumor fragments (50 mg) were implanted on the CAMs of 10 day old embryos as described in Example 5A. Twenty four hours later, embryos received a single intravenous injection of 300 ug/100 ul of either the cyclo-RADfV (69601) and or cyclo-RGDfV (66203). After a total of 72 hours, tumors were removed, examined

- 77 -

morphologically, and photographed with a stereo microscope as described in Example 9A1).

The panels shown in Figures 15A through 15E correspond as follows: Figure 15A, duplicate samples treated with cyclo-RADfV peptide (69601); Figure 15B, duplicate samples treated with cyclo-RGDfV peptide (66203); Figure 15C, adjacent CAM tissue taken from the same embryos treated with cyclo-RGDfV peptide (66203) and Figures 15D and 15E, high magnification (13x) of peptide treated tumors. Figure 15D depicts normal blood vessels from control peptide (69601) treated tumor. Figure 15E depicts examples of disrupted blood vessels from cyclo-RGDfV peptide (66203) treated tumors (arrows).

The results illustrate that only peptide 66203 inhibited vessel formation, and further that vessels in the CAM tissue adjacent to the tumor were not affected.

10. Regression of Tumor Tissue Growth With  $\alpha_v\beta_3$  Antagonists as Measured by In Vivo Rabbit Eye Model Assay

The effect of anti- $\alpha_v\beta_3$  antagonists on growth factor-induced angiogenesis can be observed in naturally transparent structures as exemplified by the cornea of the eye. New blood vessels grow from the rim of the cornea, which has a rich blood supply, toward the center of the cornea, which normally does not have a blood supply. Stimulators of angiogenesis, such as  $\beta$ FGF, when applied to the cornea induce the growth of new blood vessels from the rim of the cornea. Antagonists of angiogenesis, applied to the cornea, inhibit the growth of new blood vessels from the rim of the cornea. Thus, the cornea undergoes angiogenesis through an invasion of endothelial cells from the rim of the cornea into the tough collagen-packed



- 78 -

corneal tissue which is easily visible. The rabbit eye model assay therefore provides an in vivo model for the direct observation of stimulation and inhibition of angiogenesis following the implantation of compounds directly into the cornea of the eye.

A. In Vivo Rabbit Eye Model Assay

1) Angiogenesis Induced by Growth Factors

Angiogenesis was induced in the in vivo rabbit eye model assay with the growth factor  $\beta$ FGF and is described in the following.

a. Preparation of Hydron Pellets Containing Growth Factor and Monoclonal Antibodies

Hydron polymer pellets

containing growth factor and mAbs were prepared as described by D'Amato, et al., Proc. Natl. Acad. Sci., 91:4082-4085 (1994). The individual pellets contained 650 ng of the growth factor  $\beta$ FGF bound to sucralfate (Carafet, Marion Merrell Dow Corporation) to stabilize the  $\beta$ FGF and ensure its slow release into the surrounding tissue. In addition, hydron pellets were prepared which contained either 40  $\mu$ g of the mAb LM609 (anti- $\alpha$ v $\beta$ 3) or mAb P1F6 (anti- $\alpha$ v $\beta$ 5) in PBS. The pellets were cast in specially prepared Teflon pegs that have a 2.5 mm core drilled into their surfaces. Approximately 12  $\mu$ l of casting material was placed into each peg and polymerized overnight in a sterile hood. Pellets were then sterilized by ultraviolet irradiation.

b. Treatment with Monoclonal Antibodies

Each experiment consisted of three rabbits in which one eye received a pellet comprising  $\beta$ FGF and LM609 and the other eye

- 79 -

received a pellet comprising  $\beta$ FGF and a mouse mAb P1F6 (anti- $\alpha_v\beta_5$ ). The use of paired eye testing to compare LM609 (anti- $\alpha_v\beta_3$ ) to other mAb and PBS controls provides a means for rigorous testing to demonstrate significant differences between the mAbs tested.

The P1F6 mAb immunoreacts with the integrin  $\alpha_v\beta_5$  which is found on the surface of vascular endothelial cells but is presumably not involved in angiogenesis. To determine whether the mAb P1F6 was involved in angiogenesis, pellets containing only this mAb were prepared and assayed as described below to confirm that the mAb did not induce angiogenesis.

All of the mAbs tested were purified from ascites fluid using Protein-A Sepharose CL-4B affinity column chromatography according to well-known methods. The eluted immunoglobulin was then dialyzed against PBS and treated with Detoxi-gel (Pierce Chemicals) to remove endotoxin. Endotoxin has been shown to be a potent angiogenic and inflammatory stimulant. mAbs were therefore tested for the presence of endotoxin with the Chromogenic Limulus Amebocyte Lysate Assay (Bio-Whittaker) and only those mAbs without detectable endotoxin were used in the rabbit eye model assay.

A hydon pellet comprising  $\beta$ FGF and mAb LM609 (anti- $\alpha_v\beta_3$ ) or P1F6 (anti- $\alpha_v\beta_5$ ) was inserted into a corneal pocket formed in the eye of rabbits. The hydon pellet also contained sucralfate to stabilize the  $\beta$ FGF during the assay. Individual pellets were implanted into surgically created "pockets" formed in the mid-stroma of the cornea of rabbits. The surgical procedure was done under sterile technique using a Wild model M691 operating microscope equipped with a beamsplitter to which was mounted a camera for photographically recording individual corneas. A 3 mm by 5 mm "pocket" was

- 80 -

created in the corneal stroma by making a 3 mm incision to half the corneal thickness with a 69 Beaver blade. The stroma was dissected peripherally using an iris spatula and the pellet  
5 was implanted with its peripheral margin 2 mm from the limbus.

During the following 14 days,  $\beta$ FGF and mAb diffused from the implanted pellet into the surrounding tissue and thereby effected  
10 angiogenesis from the rim of the cornea.

Representative results of each treatment are depicted in Figures 16A through 16E. The amount of vessels present are quantitated and described in terms of clock hours which are defined as follows.  
15 The eye is divided into 12 equal sections in the same manner as a clock is divided into hours. "One clock hour of vessels" refers to that amount of vessels which fills an area of the eye equivalent to one hour on a clock. The five rabbits which  
20 received only  $\beta$ FGF exhibited florid angiogenesis in which new blood vessels had grown from the rim of the cornea toward the center of the cornea, which normally does not have blood vessels. One of these rabbits had only 1 clock hour of vessels to the  
25 pellet. Two of the rabbits which received both  $\beta$ FGF and mAb LM609 had absolutely no detectable angiogenesis up to 14 days following surgery. One of these rabbits had 3 foci of hemorrhagic and budding vessels by day 14. Two of the rabbits  
30 which received  $\beta$ FGF and mAb P3G2 (anti- $\alpha_v\beta_3$ ) showed extensive vascularization in which new blood vessels had grown from the rim of the cornea into the cornea. One of these rabbits had only 1 to 2 hours of vessels to the pellet.

35 As evidenced in the rabbit eye model assay, no angiogenic effect was observed on normal paralimbal vessels in the presence of the growth factor  $\beta$ FGF in rabbits which received mAb LM609 (anti- $\alpha_v\beta_3$ ). In

- 81 -

contrast, angiogenesis was observed on paralimbal vessels in the presence of the growth factor  $\beta$ FGF in rabbits which received the mAb P3G2 (anti- $\alpha_v\beta_3$ ). The complete inhibition of corneal angiogenesis by mAb LM609 is substantially greater than any previously reported anti-angiogenic reagent.

11. In Vivo Regression of Tumor Tissue Growth With  $\alpha_v\beta_3$  Antagonists As Measured by Chimeric Mouse:Human Assay

An in vivo chimeric mouse:human model was generated by replacing a portion of skin from a SCID mouse with human neonatal foreskin (Figure 17). After the skin graft was established, the human foreskin was inoculated with carcinoma cells. After a measurable tumor was established, either mAb LM609 (anti- $\alpha_v\beta_3$ ) or PBS was injected into the mouse tail vein. Following a 2-3 week period, the tumor was excised and analyzed by weight and histology.

A. In Vivo Chimeric Mouse:Human Assay

The in vivo chimeric mouse:human model is prepared essentially as described in Yan, et al., J. Clin. Invest., 91:986-996 (1993). Briefly, a 2 cm<sup>2</sup> square area of skin was surgically removed from a SCID mouse (6-8 weeks of age) and replaced with a human foreskin. The mouse was anesthetized and the hair removed from a 5 cm<sup>2</sup> area on each side of the lateral abdominal region by shaving. Two circular graft beds of 2 cm<sup>2</sup> were prepared by removing the full thickness of skin down to the fascia. Full thickness human skin grafts of the same size derived from human neonatal foreskin were placed onto the wound beds and sutured into place. The graft was covered with a Band-Aid which was sutured to the skin. Micropore cloth tape was also applied to cover the wound.

- 82 -

The M21L human melanoma cell line or MDA 23.1 breast carcinoma cell line (ATCC HTB 26;  $\alpha_v\beta_3$  negative by immunoreactivity of tissue sections with mAb LM609), were used to form the solid human tumors on the human skin grafts on the SCID mice. A single cell suspension of  $5 \times 10^6$  M21-L or MDA 23.1 cells was injected intradermally into the human skin graft. The mice were then observed for 2 to 4 weeks to allow growth of measurable human tumors.

#### B. Intravenous Application

##### 1) Treatment With Monoclonal Antibodies

Following the growth of measurable tumors, SCID mice, which had been injected with M21L tumor cells, were injected intravenously into the tail vein with 250  $\mu$ g of either the mAb LM609 (anti- $\alpha_v\beta_3$ ) or PBS twice a week for 2 to 3 weeks. After this time, the tumors were resected from the skin and trimmed free of surrounding tissue. Several mice were evaluated for each treatment with the average tumor weight from each treatment being calculated and shown at the bottom of Table 6.

Table 6

<u>M21L Tumor Number</u>	<u>Treatment</u>	<u>Tumor Weight (mg)</u>
1	PBS	158
2		192
3		216
4		227
5	LM609	195
6		42
7		82
8		48
9		37
10		100

11

172

	<u>Treatment</u>	<u>Average Tumor Weight (mg)</u>
	PBS	198
5	LM609	113

Exposure of the M21L  $\alpha_v\beta_3$ -negative human carcinoma tumor mass in the mouse:human chimeric assay system to LM609 (anti- $\alpha_v\beta_3$ ) caused the decrease from the PBS treated average tumor weight of 198 mg to 113 mg.

Representative examples of M21L tumors treated with the mAb LM609 (anti- $\alpha_v\beta_3$ ) and PBS were examined morphologically. The PBS-treated tumors were large (8 to 10 mm in diameter) and well vascularized whereas those treated with mAb LM609 (anti- $\alpha_v\beta_3$ ) were much smaller (3 to 4 mm in diameter) and lacked detectable blood vessels.

Tumors formed in skin grafts which had been injected with MDA 23.1 cells were detectable and measurable. Morphological examination of the established tumors revealed that neovascularization from the grafted human tissue into the MDA 23.1 tumor cells had occurred.

Thus, the blocking of the  $\alpha_v\beta_3$  receptor by the intravenous application of  $\alpha_v\beta_3$ -specific LM609 antibody resulted in a regression of a carcinoma in this model system in the same manner as the CAM and rabbit eye model systems as described in Examples 9 and 10, respectively.

#### 12. Stimulation of Vascular Cells to Enter the Cell Cycle and Undergo Apoptosis in the Presence of Antagonists of Integrin $\alpha_v\beta_3$ as Measured in the CAM Assay

The angiogenic process clearly depends on the capacity of cytokines such as  $\beta$ FGF and VEGF to stimulate vascular cell proliferation. Mignatti et

- 84 -

al., J. Cell. Biochem., 471:201 (1991); Takeshita  
et al., J. Clin. Invest., 93:662 (1994); and  
Koyama et al., J. Cell. Physiol., 158:1 (1994).  
However, it is also apparent that signaling events  
5 may regulate the differentiation of these vascular  
cells into mature blood vessels. Thus, it is  
conceivable that interfering with signals related  
to either growth or differentiation of vascular  
cells undergoing new growth or angiogenesis may  
10 result in the perturbation of angiogenesis.

Integrin ligation events have been shown to  
participate in both cell proliferation as well as  
apoptosis or programmed cell death in vitro.  
Schwartz, Cancer Res., 51:1503 (1993); Meredith et  
15 al., Mol. Biol. Cell., 4:953 (1993); Frisch et al.,  
J. Cell Biol., 124:619 (1994); and Ruoslahti et  
al., Cell, 77:477 (1994). Close examination of the  
effects of  $\alpha_v\beta_3$  antagonists on angiogenesis reveals  
the presence of discontinuous and disrupted tumor-  
20 associated blood vessels. Therefore, it is  
possible that the loss of blood vessel continuity  
may be due to selective necrosis or apoptosis of  
vascular cells.

To explore this possibility, CAMs were  
25 examined after induction of angiogenesis with the  
growth factor  $\beta$ FGF and treatment with the mAb and  
cyclic peptides of this invention.

#### A. Treatment with Monoclonal Antibodies

30 Apoptosis can be detected by a variety of  
methods which include direct examination of DNA  
isolated from tissue to detect fragmentation of the  
DNA and the detection of 3'OH in intact tissue with  
an antibody that specifically detects free 3'OH  
35 groups of fragmented DNA.

#### 1) Analysis of DNA Fragmentation

- 85 -

Angiogenesis was induced by placing filter disks saturated with  $\beta$ FGF on the CAMs of 10-day old embryos as described in Examples 6A.

Immunohistological analysis of CAMs with LM609 (anti- $\alpha_v\beta_3$ ) revealed peak expression of  $\alpha_v\beta_3$  on blood vessels 12 to 24 hrs after initiation of angiogenesis with  $\beta$ FGF. Thus, 24 hrs after stimulation with  $\beta$ FGF, embryos were inoculated intravenously with 100  $\mu$ l of PBS alone or PBS containing 300  $\mu$ g of either mAb CSAT (anti- $\beta_1$ ) or LM609 (anti- $\alpha_v\beta_3$ ).

DNA fragmentation was detected by resecting the CAM tissue directly below  $\beta$ FGF saturated filter disks 24 hr or 48 hr after intravenous inoculations with mAb LM609 (anti- $\alpha_v\beta_3$ ), CSAT (anti- $\beta_1$ ), or PBS. Resected CAM tissues were washed three times with sterile PBS and finely minced, resuspended in 0.25% bacterial collagenase (Worthington Biochemical; Freehold, NJ) and incubated for 90 minutes at 37C with occasional vortexing. DNA was extracted from equal numbers of CAM cells from single cell suspension as previously described. Bissonette, et al., Nature, 359:552 (1992). Briefly, equal numbers of CAM cells were lysed in 10 mM tris-Cl, pH 8.0, 10 mM EDTA in 0.5% (v/v) Triton X-100 (Sigma, St. Louis, MO). Cell lysates were centrifuged at 16,000x g for 15 minutes at 4C to separate soluble fragmented DNA from the intact chromatin pellet. Fragmented DNA was washed, precipitated, and analyzed on a 1.2% (w/v) agarose gel.

Soluble fragmented DNA was isolated from an equal number of CAM cells from each treatment, separated electrophoretically on an agarose gel, and visualized by staining with ethidium bromide. No difference was detected in the relative amount of DNA fragmentation resulting from the three different treatments 24 hours after treatment.



- 86 -

However, by 48 hours following treatment with mAb LM609 (anti- $\alpha_v\beta_3$ ), a significant increase in DNA fragmentation was observed when compared to embryos treated with either mAb CSAT (anti- $\beta_1$ ) or PBS alone.

5

2) Stimulation of Vascular Cells to Enter the Cell Cycle

To experimentally examine the role of  $\alpha_v\beta_3$  in these processes, cells derived from CAMs treated with or without  $\beta$ FGF were stained with propidium iodide and immunoreacted with mAb LM609 (anti- $\alpha_v\beta_3$ ).

CAMs isolated from embryos 24 and 48 hours after treatment with mAb LM609 (anti- $\alpha_v\beta_3$ ), CSAT (anti- $\beta_1$ ), or PBS were dissociated into single cell suspensions by incubation with bacterial collagenase as described above. Single cells were then permeabilized and stained with Apop Tag Insitu Detection Kit according to the manufacturer's instructions (Oncor, Gaithersburg, MD). Apop Tag is an antibody that specifically detects free 3'OH groups of fragmented DNA. Detection of such free 3'OH groups is an established method for the detection of apoptotic cells. Gavrieli et al., J. Cell Biol., 119:493 (1992).

Apop Tag stained cells were then rinsed in 0.1% (v/v) Triton X-100 in PBS and resuspended in FACS buffer containing 0.5% (w/v) BSA, 0.02% (w/v) sodium azide and 200  $\mu$ g/ml RNase A in PBS. Cells were incubated for 1.5 hrs, washed, and analyzed by fluorescence activated cell sorting. Cell fluorescence was measured using a FACScan flow cytometer and data analyzed as described below.

Cell fluorescence was measured with a FACScan flow cytometer (Becton Dickinson, Mountain View, CA). Side scatter (SSC) and forward scatter (FSC) were determined simultaneously and all data were

- 87 -

collected with a Hewlet Packard (HP9000) computer equipped with FACScan research software (Becton Dickinson, Mountain View, CA). The data were analyzed with P.C Lysis version I software (Becton Dickinson, Mountain View, CA). Negative control gates were set by using cell suspensions without the addition of primary antibodies from the Apop Tag kit. Identical gating was applied to both cell populations resulting in the analysis of approximately 8,000 cells per different cell treatment.

The percent of single cells derived from mAb treated CAMs and stained with Apop Tag as determined by FACS analysis is shown in Figure 18. The black bar represents cells from embryos treated 24 hours prior to analysis. The stippled bar represents cells from embryos treated 48 hours prior to analysis. Each bar represents the mean  $\pm$  S.E. of three replicates.

As shown in Figure 18, CAMs treated two days prior with mAb LM609 (anti- $\alpha_v\beta_3$ ) showed a 3 to 4-fold increase in Apop Tag staining as compared to CAMs treated with either PBS alone or CSAT (anti- $\beta_1$ ).

#### B. Treatment With Synthetic Peptides

CAM assays with growth factor-induced angiogenesis, as described in Example 6A, were also performed with the synthetic peptides of this invention to determine the effect of cyclic peptides on apoptosis. The peptides cyclo-RGDfV (66203) and cyclo-RADfV (69601) were prepared as described in Example 1. The peptide solutions or PBS were injected into the CAM preparation at a concentration of 300  $\mu$ g/ml. At 24 and 48 hours, the filter paper and surrounding CAM tissue was dissected and stained with the Apop Tag to detect apoptosis as described above in Example 12A2).

- 88 -

As shown in Figure 18, CAMs treated two days prior with peptide 69203 (cyclo-RGDfV) showed a 3 to 4-fold increase in Apop Tag staining as compared to CAMs treated with either PBS alone or control cyclic peptide 69601 (cyclo-RADfV).

C. Effect of Treatment With Monoclonal Antibodies on Apoptosis and Cell Cycle

Single cell suspensions were also examined for the number of copies of chromosomal DNA by staining with propidium iodide to determine the effect of treatment with monoclonal antibodies on the cell cycle and for apoptosis by staining with the Apop Tag.

Single cell suspensions of CAMS treated 24 or 48 hours prior with mAb LM609 (anti- $\alpha_v\beta_3$ ) or CSAT (anti- $\beta_1$ ) or PBS were prepared as described in Example 12A1).

For staining of cells with the Apop Tag, cell suspensions were washed three times with buffer containing 2.5% (w/v) BSA and 0.25% (w/v) sodium azide in PBS. Cells were then fixed in 1% (w/v) paraformaldehyde in PBS for 15 minutes followed by three washes as described above. To prevent nonspecific binding, single cell suspensions were blocked with 5% (w/v) BSA in PBS overnight at 4C. Cells were then washed as before, stained with Apop Tag, and cell fluorescence measured with a FACScan as described above in Example 12A.

Cells from each experimental condition were stained with propidium iodide (Sigma, St. Louis, MO) at 10  $\mu$ g/ml in PBS for 1 hour, washed two times with PBS, and analyzed for nuclear characteristics typical of apoptosis, including chromatin condensation and segmentation. The percentage of apoptotic cells were estimated by morphological analysis of cells from at least 10 to 15 randomly selected microscopic fields.

- 89 -

The combined results of single cell suspensions of CAMs from embryos treated with either CSAT (anti- $\beta_1$ ) or LM609 (anti- $\alpha_v\beta_3$ ), stained with Apop Tag and propidium iodide, and analyzed by FACS are given in Figure 19. The Y axis represents Apop Tag staining (apoptosis), the X axis represents propidium iodide staining (DNA content). The horizontal line represents the negative gate for Apop Tag staining. The left and right panels indicate CAM cells from CSAT and LM609 treated embryos, respectively. Cell cycle analysis was performed by analysis of approximately 8,000 events per condition and data represented in a contour plot.

Samples of single cells stained with the DNA dye propidium iodide revealed that 25-30% of the LM609 (anti- $\alpha_v\beta_3$ ) treated CAM cells 48 hours after treatment showed evidence of nuclear condensation and/or segmentation. These processes are characteristic of cells undergoing apoptosis. This is in contrast to CAMs treated with CSAT (anti- $\beta_1$ ) where 90-95% of the cells showed normal nuclear staining.

As shown in Figure 19, consistent with the induction of apoptosis by LM609, a significant number of cells in a peak containing less than one copy of DNA was observed (AO). This peak has been previously shown to represent fragmented DNA in late stage apoptotic cells. Telford et al., Cytometry, 13:137 (1992). Furthermore, these AO cells readily stain with Apop Tag confirming the ability of this reagent to detect apoptotic cells. However, in addition to the staining of cells in AO, a significant number of cells containing greater than one copy of DNA also stained with Apop Tag (Figure 19). These results demonstrate the LM609 has the ability to promote apoptosis among vascular cells that had already entered the cell

- 90 -

cycle. In contrast, cells derived from control CAMs which had entered the cell cycle showed minimal Apop Tag staining consistent with the few apoptotic cells detected in control treated CAMs.

5           Among those cells in the  $\beta$ FGF stimulated CAMs that had entered the cell cycle (S and G2/M phase), 70% showed positive staining with LM609 (anti- $\alpha_v\beta_3$ ). This is compared to 10% LM609 staining observed among cycling cells from non- $\beta$ FGF treated CAMs.

10          These findings indicate that after  $\beta$ FGF stimulation, the majority of the  $\alpha_v\beta_3$ -bearing cells show active proliferation.

          Taken together these findings indicate that intravenous injection of mAb LM609 or cyclic peptide antagonist of  $\alpha_v\beta_3$  promote apoptosis within

15          the chick CAM following induction of angiogenesis.

          CAMs were also examined histologically for expression of  $\alpha_v\beta_3$  by immunoreactivity with LM609 and for cells which were undergoing apoptosis by

20          immunoreactivity with Apop Tag. CAM sections resected from embryos treated 48 hrs prior with LM609 (anti- $\alpha_v\beta_3$ ), CSAT (anti- $\beta_1$ ), or PBS prepared in Example 5A were washed, embedded in OTC (Baxter) and snap frozen in liquid nitrogen. Six micron

25          sections of CAM tissues were cut, fixed in acetone for 30 seconds, and stored at -70C until use. Tissue sections were prepared for staining by a brief rinse in 70% (v/v) ethanol (ETOH) followed by washing three times in PBS. Next, sections were

30          blocked with 5% (w/v) BSA in PBS for 2 hrs, followed by incubation with 10  $\mu$ g/ml of mAb LM609 for 2 hrs. The sections were then washed and incubated with a 1:50 dilution of rhodamine conjugated goat anti-mouse IgG (Fisher Scientific, Pittsburgh, PA) for 2 hrs. Finally, the same

35          sections were washed and stained with the Apop Tag as described in Example 12A2). The stained tissue

- 91 -

sections were mounted and analyzed by confocal immunofluorescent microscopy.

In Figure 20, panels A through C represent CAM tissue from CSAT (anti- $\beta_1$ ) treated embryos and panels D through F represent CAM tissue from LM609 (anti- $\alpha_v\beta_3$ ) treated embryos. Panels A and D depict tissues stained with Apop Tag and visualized by fluorescence (FITC) superimposed on a D.I.C. image. Panels B and E depict the same tissues stained with mAb LM609 (anti- $\alpha_v\beta_3$ ) and visualized by fluorescence (rhodamine). Panels C and F represent merged images of the same tissues stained with both Apop Tag and LM609 where yellow staining represents colocalization. The bar represents 15 and 50  $\mu\text{m}$  in the left and right panels, respectively.

As shown in Figure 20 (A-C), after intravenous injection of CSAT or PBS control, staining with Apop Tag appeared minimal and random, indicating a minimal level of apoptosis within the tissue. In contrast, CAMs from embryos previously treated with LM609 or cyclic peptide 203 showed a majority of the vessels staining intensely with Apop Tag while minimal reactivity was observed among surrounding nonvascular cells (Figure 20 D-F). Furthermore, when both Apop Tag and LM609 were used to stain these tissues (19C and 19F) significant colocalization was only observed between these markers in CAMs derived from embryos treated with  $\alpha_v\beta_3$  antagonists (Figure 20F). These findings demonstrate that after induction of angiogenesis in vivo, inhibitors of integrin  $\alpha_v\beta_3$  selectively promote apoptosis of  $\alpha_v\beta_3$ -bearing blood vessels.

While angiogenesis is a complex process involving many molecular and cell biological events, several lines of evidence suggest that vascular cell integrin  $\alpha_v\beta_3$  plays a relatively late role in this process. First, immunohistological

- 92 -

analysis reveals that expression of  $\alpha_v\beta_3$  on vascular cells reached a maximum 12-24 hours after the induction of angiogenesis with  $\beta$ FGF. Secondly, antagonists of  $\alpha_v\beta_3$  perturb angiogenesis induced by multiple activators suggesting that this receptor is involved in common pathway downstream from perhaps all primary signaling events leading to angiogenesis. Thirdly, mAb LM609 or cyclic peptide treated CAMs did not show a significant increase in apoptosis as measured by DNA laddering until 48 hours post treatment with these antagonists. Finally, antagonists of  $\alpha_v\beta_3$  promote apoptosis of vascular cells that have already been induced to enter the cell cycle.

The results presented herein provide the first direct evidence that integrin ligation events can regulate cell survival in vivo. It is therefore hypothesized that once angiogenesis begins, individual vascular cells divide and begin to move toward the angiogenic source, after which,  $\alpha_v\beta_3$  ligation provides a signal allowing continued cell survival which leads to differentiation and the formation of mature blood vessels. However, if  $\alpha_v\beta_3$  ligation is prevented then the cells fail to receive this molecular cue and the cells go into apoptosis by default. This hypothesis would also predict that after differentiation has occurred mature blood vessels no longer require  $\alpha_v\beta_3$  signaling for survival and thus are refractory to antagonists of this integrin.

Finally, the results presented herein provide evidence that antagonists of integrin  $\alpha_v\beta_3$  may provide a powerful therapeutic approach for the treatment of neoplasia or other diseases characterized by angiogenesis. First, antagonists of  $\alpha_v\beta_3$  disrupt newly forming blood vessels without affecting the pre-existing vasculature. Second, these antagonists had no significant effect on

- 93 -

chick embryo viability, suggesting they are non-toxic. Third, angiogenesis was significantly blocked regardless of the angiogenic stimuli. Finally, systemic administration of  $\alpha_v\beta_3$  antagonists causes dramatic regression of various histologically distinct human tumors.

Thus, the aforementioned Examples demonstrate that integrin  $\alpha_v\beta_3$  plays a key role in angiogenesis induced by a variety of stimuli and as such  $\alpha_v\beta_3$  is a valuable therapeutic target with the  $\alpha_v\beta_3$  antagonists of this invention for diseases characterized by neovascularization.

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the cell line deposited, since the deposited embodiment is intended as a single illustration of one aspect of the invention and any cell line that is functionally equivalent is within the scope of this invention. The deposit of material does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustration that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.



- 94 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: The Scripps Research Institute
- (B) STREET: 10666 North Torrey Pines Road
- (C) CITY: La Jolla
- (D) STATE: CA
- (E) COUNTRY: USA
- (F) POSTAL CODE (ZIP): 92037
- (G) TELEPHONE: 619-554-2937
- (H) TELEFAX: 619-554-6312

(ii) TITLE OF INVENTION: METHODS AND COMPOSITIONS USEFUL FOR  
INHIBITION OF ANGIOGENESIS

(iii) NUMBER OF SEQUENCES: 14

## (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

## (v) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER: PCT/US 95/
- (B) FILING DATE: 09-MAR-1995

## (vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 08/210,715
- (B) FILING DATE: 18-MAR-1994

## (vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 08/366,665
- (B) FILING DATE: 30-DEC-1994

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

- 95 -

## (ix) FEATURE:

- (A) NAME/KEY: Peptid
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /label= BOC-GRGDFV-OMe  
/note= "BOC signifies the N-terminal protecting group butyloxycarb nyl; OMe signifies a C-terminal methyl ester; arginine in the second position"

## (ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /label= OMe  
/note= "OMe signifies the C-terminal protecting group methyl ester."

## (ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /label= D-Arg  
/note= "A prefix "D" in D-Arg signifies that the arginine in position 2 is a D-amino acid."

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Gly Arg Gly Asp Phe Val  
1 5

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## (v) FRAGMENT TYPE: internal

## (ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /label= BOC  
/note= "BOC signifies the N-terminal blocking group tertbutyloxycarbonyl."

## (ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /label= OH

- 96 -

/note= "OH signifies a free C-terminal carboxylic acid."

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /label= D-Arg  
/note= "A prefix "D" in D-Arg signifies that the arginine in position 2 is a D-amino acid."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Gly Arg Gly Asp Phe Val  
1 5

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 6 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /label= H  
/note= "H signifies a free N-terminal amine."

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /label= OH  
/note= "OH signifies a free C-terminal carboxylic acid."

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /label= D-Arg  
/note= "A prefix "D" in D-Arg at position 2, signifies that the arginine is a D-amino acid."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

- 97 -

Gly Arg Gly Asp Phe Val  
1 5

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

## (ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /label= cyclo  
/note= "Cyclo signifies a cyclic peptide; lower  
case letters indicate a D-amino acid; capital  
letters indicate a L-amino acid."

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Gly Arg Gly Asp Phe Val  
1 5

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

## (ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..5
- (D) OTHER INFORMATION: /label= cyclo  
/note= "Cyclo signifies a cyclic peptide; l wer

- 98 -

case letters indicate a D-amino acid; capital letters indicate a L-amino acid."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Arg Gly Asp Phe Val  
1 5

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..5
- (D) OTHER INFORMATION: /label= cyclo  
/note= "Cyclo signifies a cyclic peptide; lower case letters indicate a D-amino acid; capital letters indicate a L-amino acid."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Arg Ala Asp Phe Val  
1 5

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

- 99 -

## (ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..5
- (D) OTHER INFORMATION: /label= cyclo  
/note= "Cyclo signifies a cyclic peptide; lower  
case letters indicate a D-amino acid; capital  
letters indicate a L-amino acid."

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Arg Gly Asp Phe Val  
1 5

## (2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 15 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Tyr Thr Ala Glu Cys Lys Pro Gln Val Thr Arg Gly Asp Val Phe  
1 5 10 15

## (2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 5 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

## (ix) FEATURE:

- (A) NAME/KEY: Peptide

- 100 -

- (B) LOCATION: 1..5
- (D) OTHER INFORMATION: /label= cyclo  
/note= "Cyclo signifies a cyclic peptide; lower  
case letters indicate a D-amino acid; capital  
letters indicate a L-amino acid."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Arg Ala Asp Phe Val  
1 5

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 6 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..6
- (D) OTHER INFORMATION: /label= cyclo  
/note= "Cyclo signifies a cyclic peptide; lower  
case letters indicate a D-amino acid; capital  
letters indicate a L-amino acid."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Ala Arg Gly Asp Phe Leu  
1 5

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 6 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- 101 -

(v) FRAGMENT TYPE: internal

(ix) FEATURE:

(A) NAME/KEY: Peptide

(B) LOCATION: 1..6

(D) OTHER INFORMATION: /label= cyclo

/note= "Cyclo signifies a cyclic peptide; lower  
case letters indicate a D-amino acid; capital  
letters indicate a L-amino acid."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Gly Arg Gly Asp Phe Leu  
1 5

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Thr Arg Gln Val Val Cys Asp Leu Gly Asn Pro Met  
1 5 10

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal



- 102 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Gly	Val	Val	Arg	Asn	Asn	Glu	Ala	Leu	Ala	Arg	Leu	Ser
1				5				10				

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Thr	Asp	Val	Asn	Gly	Asp	Gly	Arg	His	Asp	Leu
1			5					10		

- 103 -

What Is Claimed Is:

1. A method for inhibiting angiogenesis in a tissue comprising administering to said tissue a composition comprising an angiogenesis-inhibiting amount of an  $\alpha_v\beta_3$  antagonist.  
5
2. The method of claim 1 wherein said  $\alpha_v\beta_3$  antagonist inhibits binding of fibrinogen to  $\alpha_v\beta_3$  but does not substantially inhibit binding of fibrinogen to  $\alpha_{Ib}\beta_3$  or  $\alpha_v\beta_1$ .
3. The method of claim 1 wherein said  $\alpha_v\beta_3$  antagonist is a monoclonal antibody immunospecific for  $\alpha_v\beta_3$ .  
10
4. The method of claim 3 wherein said monoclonal antibody has the immunoreaction characteristics of the monoclonal antibody LM609 (ATCC HB 9537).  
15
5. The method of claim 1 wherein said  $\alpha_v\beta_3$  antagonist is an RGD-containing polypeptide.
6. The method of claim 5 wherein said polypeptide is selected from the group consisting of c-(GrGDFV) (SEQ ID NO 4), c-(RGDFv) (SEQ ID NO 5), c-(RGDFv) (SEQ ID NO 7), and YTAECKPQVTRGDVF (SEQ ID NO 8), and a salt thereof.  
20
7. The method of claim 6 wherein said salt is hydrochloride or trifluoroacetate.  
25
8. The method of claim 1 wherein said tissue is inflamed and said angiogenesis is inflamed tissue angiogenesis.
9. The method of claim 8 wherein said tissue is arthritic.  
30
10. The method of claim 9 wherein said arthritic tissue is present in a mammal with rheumatoid arthritis.
11. The method of claim 1 wherein said tissue is the retinal tissue of a patient with diabetic retinopathy and said angiogenesis is retinal angiogenesis.  
35

- 104 -

12. The method of claim 1 wherein said tissue is a hemangioma.

13. The method of claim 1 wherein said tissue is a solid tumor or a solid tumor metastasis and said angiogenesis is tumor angiogenesis.

14. The method of claim 1 wherein said angiogenesis-inhibiting amount is from about 2  $\mu\text{M}$  to 5 mM.

15. The method of claim 1 wherein said administering comprises intravenous, transdermal, intrasynovial, intramuscular, or oral administration.

16. The method of claim 1 wherein said administering is conducted in conjunction with chemotherapy.

17. The method of claim 1 wherein said administering comprises a single dose intravenously.

18. A method for regression of established tumor in a tissue comprising administering to said tissue a composition comprising an angiogenesis-inhibiting amount of an  $\alpha_v\beta_3$  antagonist.

19. The method of claim 18 wherein said  $\alpha_v\beta_3$  antagonist inhibits binding of fibrinogen to  $\alpha_v\beta_3$  but does not substantially inhibit binding of fibrinogen to  $\alpha_{IIb}\beta_3$  or  $\alpha_v\beta_1$ .

20. The method of claim 18 wherein said  $\alpha_v\beta_3$  antagonist is a monoclonal antibody immunospecific for  $\alpha_v\beta_3$ .

21. The method of claim 20 wherein said monoclonal antibody has the immunoreaction characteristics of the monoclonal antibody LM609 (ATCC HB 9537).

22. The method of claim 18 wherein said  $\alpha_v\beta_3$  antagonist is an RGD-containing polypeptide.

23. The method of claim 22 wherein said polypeptide is selected from the group consisting of c-(GrGDFV) (SEQ ID NO 4), c-(RGDfV) (SEQ ID NO

- 105 -

5), c-(RGDFv) (SEQ ID NO 7), and YTAECKPQVTRGDVF (SEQ ID NO 8), and a salt thereof.

24. The method of claim 23 wherein said salt is hydrochloride or trifluoroacetate.

5           25. The method of claim 18 wherein said tissue is inflamed and said angiogenesis is inflamed tissue angiogenesis

26. The method of claim 25 wherein said tissue is arthritic.

10           27. The method of claim 26 wherein said arthritic tissue is present in a mammal with rheumatoid arthritis.

28. The method of claim 18 wherein said tissue is the retinal tissue of a patient with  
15           diabetic retinopathy and said angiogenesis is retinal angiogenesis.

29. The method of claim 18 wherein said tissue is a hemangioma.

30. The method of claim 18 wherein said  
20           tissue is a solid tumor or a solid tumor metastasis and said angiogenesis is tumor angiogenesis.

31. The method of claim 18 wherein said angiogenesis-inhibiting amount is from about 2  $\mu$ M to 5 mM.

25           32. The method of claim 18 wherein said administering comprises intravenous, transdermal, intrasynovial, intramuscular, or oral administration.

33. The method of claim 18 wherein said  
30           administering is conducted in conjunction with chemotherapy.

34. The method of claim 18 wherein said administering comprises a single dose intravenously.

35           35. The method of claim 18 wherein said regression occurs 7 days following administration of said angiogenesis-inhibiting antagonist.

- 106 -

36. A method of inducing apoptosis in neo-vasculature in a tissue comprising administering to said tissue a composition comprising an angiogenesis-inhibiting amount of an  $\alpha_v\beta_3$  antagonist.

37. The method of claim 36 wherein said  $\alpha_v\beta_3$  antagonist inhibits binding of fibrinogen to  $\alpha_v\beta_3$  but does not substantially inhibit binding of fibrinogen to  $\alpha_{IIb}\beta_3$  or  $\alpha_v\beta_1$ .

38. The method of claim 36 wherein said  $\alpha_v\beta_3$  antagonist is a monoclonal antibody immunospecific for  $\alpha_v\beta_3$ .

39. The method of claim 38 wherein said monoclonal antibody has the immunoreaction characteristics of the monoclonal antibody LM609 (ATCC HB 9537).

40. The method of claim 36 wherein said  $\alpha_v\beta_3$  antagonist is an RGD-containing polypeptide.

41. The method of claim 40 wherein said polypeptide is selected from the group consisting of c-(GrGDFV) (SEQ ID NO 4), c-(RGDfV) (SEQ ID NO 5), c-(RGDFv) (SEQ ID NO 7), and YTAECKPQVTRGDVF (SEQ ID NO 8), and a salt thereof.

42. The method of claim 41 wherein said salt is hydrochloride or trifluoroacetate.

43. The method of claim 36 wherein said tissue is inflamed and said angiogenesis is inflamed tissue angiogenesis.

44. The method of claim 43 wherein said tissue is arthritic.

45. The method of claim 44 wherein said arthritic tissue is present in a mammal with rheumatoid arthritis.

46. The method of claim 36 wherein said tissue is the retinal tissue of a patient with diabetic retinopathy and said angiogenesis is retinal angiogenesis.

- 107 -

47. The method of claim 36 wherein said tissue is a hemangioma.

48. The method of claim 36 wherein said tissue is a solid tumor or a solid tumor metastasis and said angiogenesis is tumor angiogenesis.

49. The method of claim 36 wherein said angiogenesis-inhibiting amount is from about 2  $\mu$ M to 5 mM.

50. The method of claim 36 wherein said administering comprises intravenous, transdermal, intrasynovial, intramuscular, or oral administration.

51. The method of claim 36 wherein said administering is conducted in conjunction with chemotherapy.

52. The method of claim 36 wherein said administering comprises a single dose intravenously.

53. The method of claim 36 wherein said apoptosis occurs at least 48 hours following administration of said angiogenesis-inhibiting antagonist.

54. The method of claim 1 wherein said tissue is a coronary artery at risk of restenosis following coronary angioplasty.

**Anti- $\beta$ 3**



**Normal Skin**

**FIG. 1A**



**Granulation Tissue**

**FIG. 1B**

**Anti- $\beta$ 3**

**Anti- $\beta$ 1**



**Normal Skin**

**FIG. 1C**



**Granulation Tissue**

**FIG. 1D**

**Anti- $\beta$ 1**

2/23

**Normal Skin**


**Anti-vWF**



**FIG. 2A**

**Granulation Tissue**

**Anti-vWF**




**FIG. 2B**



3/23

**Normal Skin**


**Anti-laminin**



**FIG. 2C**


**Granulation Tissue**

**Anti-laminin**



**FIG. 2D**

4/23



**Bladder Cancer**


**FIG. 3A**



**Colon Cancer**


**FIG. 3B**

5/23



**Breast Cancer**

**FIG. 3C**




**Lung Cancer**

**FIG. 3D**



6/23

**Control**



**FIG. 4**

7/23

**FIG. 5A****Untreated  
Anti- $\beta_1$** **FIG. 5B****Untreated  
Anti- $\alpha\beta_3$** **FIG. 5C****bFGF Treated  
Anti- $\alpha\beta_3$** 

8/23




FIG. 6

9/23

TNF $\alpha$




FIG. 7C

bFGF




FIG. 7B

Control




FIG. 7A

10/23

**FIG. 8A**  
**Control**



**bFGF**



**FIG. 8B**



**FIG. 8C**  
**Anti-β<sub>1</sub>**



**FIG. 8D**  
**Anti-α<sub>v</sub>β<sub>5</sub>**



**FIG. 8E**  
**Anti-α<sub>v</sub>β<sub>3</sub>**



11/23

**FIG. 9C**




**Cyclic RGD  
Adjacent CAM**

**FIG. 9B**



**Cyclic RGD  
Tumor**

**FIG. 9A**



**Control Peptide  
Tumor**

12/23

**FIG. 10C**  
**Anti- $\alpha$ V $\beta$ <sub>3</sub> (LM609)**



**FIG. 10B**  
**Anti- $\alpha$ V $\beta$ <sub>5</sub> (P3G2)**



**FIG. 10A**  
**Control**




RECTIFIED SHEET (RULE 91)




**FIG. 11A**

**Anti- $\beta$ 1**



**FIG. 11B**


**Anti- $\alpha$ v $\beta$ 5**



**FIG. 11C**

**Anti- $\alpha$ v $\beta$ 3**

14/23

**FIG. 12**

15/23




FIG. 13B




FIG. 13D




FIG. 13A





FIG. 13C


16/23

**FIG. 14A**




**Anti- $\alpha$ V $\beta$ <sub>5</sub>  
(P3G2)**

**FIG. 14B**




**Anti- $\alpha$ V $\beta$ <sub>3</sub>  
(LM609)**

17/23

**FIG. 15A****FIG. 15B****FIG. 15C****Control Peptide Tumor****Cyclic RGD Tumor****Cyclic RGD  
Adjacent CAM****FIG. 15D****Control Peptide****FIG. 15E****Cyclic RGD Peptide**

18/23

**P1F6**  
**(Anti- $\alpha_v\beta_5$ )**




**FIG. 16A**

**LM609**  
**(Anti- $\alpha_v\beta_3$ )**

**FIG. 16C****FIG. 16D****FIG. 16B****FIG. 16E**



19/23

**Transplant  
Human Skin****Heal 4 Weeks****Injection  
of Human  
Tumor Cell****Tumor Growth 4 Weeks****FIG. 17**

20/23




FIG. 18

21/23




FIG. 19

RECTIFIED SHEET (RULE 91)


22/23



**FIG. 20A**



**FIG. 20B**




**FIG. 20C**

RECTIFIED SHEET (RULE 91)


23/23



**FIG. 20D**



**FIG. 20E**



**FIG. 20F**

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/03035

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :A61K 39/395; C07K 16/00, 16/18, 16/28, 16/30

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/143.1, 145.1, 158.1, 155.1, 174.1; 530/388.22, 388.24, 388.25, 388.8

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EMBASE, BIOSIS, WPIDS, MEDLINE, BITECHDS, DISSABS, CONFSCI, LIFESCI, SCISEARCH, JAPIO, JICST-EPLUS, APS

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ANTIBODY, IMMUNOCONJUGATES, AND RADIOPHARMACEUTICALS, Volume 4, Number 2, Issued 1991, Mueller et al., "Pre-Clinical Therapy of Human Melanoma with Morpholino-Doxorubicin Conjugated to a Monoclonal Antibody Directed Against an Integrin on Melanoma Cells", pages 99-106, see entire document.	16, 33, 51
Y	JOURNAL OF CLINICAL INVESTIGATION, Volume 90, Issued October 1992, Nip et al. "Human Melanoma Cells Derived from Lymphatic Metastases Use Integrin $\alpha_v\beta_3$ to Adhere to Lymph Node Vitronectin", pages 1406-1413, see entire document.	1-4, 8-21, 25-39, 43-53
Y	US, A, 5,112,946 (THEODORE MAIONE) 12 MAY 1992, See entire document.	1-4, 8-21, 25-39, 43-53

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" Later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

03 JUNE 1995

Date of mailing of the international search report

16 JUN 1995

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SUSAN A. LORING

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/03035

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 5,262,520 (PLOW ET AL) 16 November 1993, see entire document.	1-4, 8-21, 25-39, 43-53
X	THE EMBO JOURNAL, Volume 4, Number 10, Issued 1985, Suzuki et al, "Complete Amino Acid Sequence of Human Vitronectin Deduced from cDNA. Similarity of Cell Attachment Sites in Vitronectin and Fibronectin", pages 2519-2524, see especially abstract and Figure 4.	5-7, 22-24, 40-42
Y	WO, A, 89/05155 (DAVID A. CHERESH) 15 JUNE 1989, see entire document.	1-54
Y	EP, A2, 0,485,749 (WOLFGANG A. WRASIDIO) 20 MAY 1992, See entire document.	16, 33, 51
Y	WO, A, 90/14103 (DAVID A. CHERESH) 29 NOVEMBER 1990, See entire document.	1-54
P, Y	THE JOURNAL OF BIOLOGICAL CHEMISTRY, Volume 269, Number 32, Issued 12 August 1994, Pfaff et al, "Selective Recognition of Cyclic RGD Peptides of NMR Defined Conformation by $\alpha$ IIb $\beta$ 3, $\alpha$ V $\beta$ 3, and $\alpha$ 5 $\beta$ 1 Integrins", pages 20233-20238, see entire document.	1-54
Y	EXPERIMENTAL HEMATOLOGY, Volume 20, Issued 1992, Leven et al, "Extracellular Matrix Stimulation of Guinea Pig Megakaryocyte Proplatelet Formation in vitro Is Mediated Through the Vitronectin Receptor", pages 1316-1322, see entire document.	1-54
X	CANCER RESEARCH, Volume 52, Issued 15 April 1992, Lafrenie et al, "Up-regulated Biosynthesis and Expression of Endothelial Cell Vitronectin Receptor Enhances Cancer Cell Adhesion", pages 2202-2208, see entire document.	1-54

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/03035

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/143.1, 145.1, 158.1, 155.1, 174.1; 530/388.22, 388.24, 388.25, 388.8